



Prognosis in Severe Brain Injury

Robert D. Stevens, MD¹⁻⁴; Raoul Sutter, MD¹⁻³

Background: The prediction of neurologic outcome is a fundamental concern in the resuscitation of patients with severe brain injury.

Objective: To provide an evidence-based update on neurologic prognosis following traumatic brain injury and hypoxic-ischemic encephalopathy after cardiac arrest.

Data Source: Search of the PubMed database and manual review of bibliographies from selected articles to identify original data relating to prognostic methods and outcome prediction models in patients with neurologic trauma or hypoxic-ischemic encephalopathy.

Data Synthesis and Conclusion: Articles were scrutinized regarding study design, population evaluated, interventions, outcomes, and limitations. Outcome prediction in severe brain injury is reliant on features of the neurologic examination, anatomical and physiological changes identified with CT and MRI, abnormalities detected with electroencephalography and evoked potentials, and physiological and biochemical derangements at both

the brain and systemic levels. Use of such information in univariable association studies generally lacks specificity in classifying neurologic outcome. Furthermore, the accuracy of established prognostic classifiers may be affected by the introduction of outcome-modifying interventions, such as therapeutic hypothermia following cardiac arrest. Although greater specificity may be achieved with scoring systems derived from multivariable models, they generally fail to predict outcome with sufficient accuracy to be meaningful at the single patient level. Discriminative models which integrate knowledge of genetic determinants and biologic processes governing both injury and repair and account for the effects of resuscitative and rehabilitative care are needed. (*Crit Care Med* 2013; 41:1104–1123)

Key Words: cardiac arrest; coma; Corticosteroid Randomization After Significant Head Injury; hypoxic-ischemic encephalopathy; International Mission on Prognosis and Analysis of Clinical Trials in Traumatic Brain Injury; prognosis; traumatic brain injury

BURDEN OF SEVERE BRAIN INJURY

The resuscitation of patients with severe brain injury (SBI), which we define as an insult severe enough to cause an acute and persistent loss of consciousness and to entail a significant likelihood of death or of long-term disability, is a central concern in intensive care medicine. Although the etiologies of SBI are diverse, the most prevalent causes are traumatic brain injury (TBI) and cardiac arrest (CA). Severe TBI is the leading cause

of death during the first five decades of life, with more than 50,000 deaths annually and functional disability in more than 40% of survivors in the United States (1). The prevalence of CA is estimated at half a million cases per year in the United States with less than 30% of patients surviving until hospital discharge and high rates of disability in survivors despite advances in cardiopulmonary resuscitation and post-CA care (2–5).

The introduction of therapeutic hypothermia for comatose survivors after CA has improved not only survival but also the functional status of survivors (4–7). Randomized trials have found that up to 50% of comatose CA patients treated with hypothermia may have a good 1-year neurologic outcome (4, 5, 8). With declining mortality, there is a pressing need to accurately and reliably predict the likelihood of functional recovery, yet existing prognostic models have many limitations. They do not recognize potentially critical determinants of outcome, such as genetic susceptibility, biological heterogeneity, and the effects of incremental refinements in intensive care and rehabilitation. Current prognostic models in SBI generally estimate the probability of a dichotomous endpoint, usually an unfavorable outcome defined as death or severe disability; however, there is intrinsic interest in understanding the likelihood of a favorable outcome, or in being able to discriminate individual probabilities in a range of outcomes, not just binary favorable/

¹Division of Neurosciences Critical Care, Department of Anesthesiology and Critical Care Medicine, Johns Hopkins University School of Medicine, Baltimore, MD.

²Division of Neurosciences Critical Care, Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, MD.

³Division of Neurosciences Critical Care, Department of Neurosurgery, Johns Hopkins University School of Medicine, Baltimore, MD.

⁴Division of Neurosciences Critical Care, Department of Radiology, Johns Hopkins University School of Medicine, Baltimore, MD.

This work was performed at Johns Hopkins University School of Medicine. Dr. Stevens has received grant support from the National Institutes of Health. Dr. Sutter has received grant support from the Research Fund of the University of Basel, The Scientific Society Basel, and the Gottfried Julia Bangerter-Rhyner Foundation.

For information regarding this article, E-mail: rstevens@jhmi.edu

Copyright © 2013 by the Society of Critical Care Medicine and Lippincott Williams & Wilkins

DOI: 10.1097/CCM.0b013e318287ee79

unfavorable ones. Finally, the unfounded assumption of an unfavorable prognosis may become “self-fulfilling” if life-sustaining treatment is withheld on the basis of that belief (9). An ideal prognostic system would help clinicians align realistic estimates of recovery with the mobilization of therapeutic resources at the individual level, thus defeating self-fulfilling prophecies. Consideration of these factors is likely to increase the relevance and accuracy of prognostic algorithms in SBI.

RECOVERY FOLLOWING SBI

The clinical trajectory following SBI can be considered in terms of three basic processes: 1) the emergence of conscious awareness, 2) the recovery of higher neuropsychological processing, and 3) the return of functional capacity (10). States commonly seen following coma are the vegetative state (VS) (11, 12), the minimally conscious state (MCS) (13), and emergence from minimally conscious state (EMCS), marked by reliable functional communication and functional object use after MCS (14); further stages of recovery are characterized by the restoration of higher cognitive functions, such as attention, memory, and executive functions (Fig. 1). The time course of recovery is variable; in a subset of patients, phenotypic characteristics may remain fixed (“plateau”), resulting in states of chronically impaired consciousness.

In contrast to coma, patients in a VS demonstrate signs of arousal; however, they are incapable of self or contextual awareness (15). The pattern of brain damage can be substantially different in VS patients depending on the etiology. A common neuropathologic finding after CA is diffuse or multifocal cortical ischemia, often associated with lesions in the thalamus and basal ganglia injury (16), with relative sparing of the brainstem (17). In TBI, the most frequent pattern is

traumatic axonal injury with lesions clustering in the corpus callosum and brainstem (18). Clinical descriptors of VS may not accurately identify underlying conscious processes; with the help of functional neuroimaging or high-density electroencephalography (EEG), it has been demonstrated that a subset of patients who meet criteria for VS have patterns of cortical activation that are analogous to those of conscious controls (19, 20). MCS is also generally seen in patients with extensive hemispheric damage, although cortico-cortical and corticothalamic connectivity may be better preserved in MCS than in VS (21–23). In contrast to VS, patients with MCS present with unequivocal signs of self or environmental awareness; however, these behaviors are not demonstrated in a reliable fashion (13).

The natural history of patients who are in VS and MCS is not well studied. In a retrospective analysis, 50% of MCS patients and only 3% of VS patients had no or moderate disabilities when evaluated 1 year after injury, and improvement was more rapid and pronounced in the patients who had a traumatic etiology (24). The appearance of visual pursuit may precede recovery of more distinct signs of consciousness (25), and an association has been observed between preserved visual pursuit and the likelihood of functional recovery (26, 27). The time course of recovery may be prognostically significant. In a multicenter study of VS and MCS patients admitted to rehabilitation centers, prediction of functional outcome correlated with the rate of improvement measured with repeated administration of the Disability Rating Scale over the first 2 weeks of observation (28). After recovery of awareness, patients recovering from SBI may have significant neurological impairments, including motor deficits, myoclonus, dystonia, movement disorders, aphasia, neglect, and abulia (29–32). In addition, many survivors of SBI have impairments in attention, memory, executive control, mood disorders,

	Brain death	Coma	Vegetative state	Minimally conscious state	Emergence from minimally conscious state
Eye opening	none	none	occasional	Occasional but inconsistent	present
Visual fixation or pursuit	none	none	none	Occasional but inconsistent	present
Spontaneous movement	none	none	nonpurposeful	present	purposeful
Response to noxious stimuli	none	none, flexion or extension	variable	variable	variable
Verbalization	none	none	none	Occasional but inconsistent	present
Functional communication	none	none	none	Occasional but inconsistent	consistent

Figure 1. Recovery phenotypes following severe brain injury.

TABLE 1. Prognostic Variables Included in the International Mission on Prognosis and Analysis of Clinical Trials and Adapted Corticoid Randomization After Significant Head Injury Models

International Mission on Prognosis and Analysis of Clinical Trials		
Basic Model	Core Model	Lab Model
Age	Age	Age
GCS	GCS	GCS
Motor score	Motor score	
Pupillary reactivity	Pupillary reactivity	Pupillary reactivity
	Marshall CT classification	Marshall CT classification
	Epidural hematoma	Epidural hematoma
	Traumatic subarachnoid hemorrhage	Traumatic subarachnoid hemorrhage
	Hypoxia	Hypoxia
	Hypotension	Hypotension
		Blood glucose level
		Hemoglobin level
Corticoid Randomization After Significant Head Injury		
Basic Model	Extended Model	
Age	Age	
GCS	GCS	
Motor score	Motor score	
Pupillary reactivity	Pupillary reactivity	
Major extracranial injury	Major extracranial injury	
	Marshall CT classification	
	Traumatic subarachnoid hemorrhage	

GCS = Glasgow Coma Scale.
Adapted from Roozenbeek et al (36).

and seizure disorders that have a dramatic impact on the level of functional independence and quality of life (33).

PROGNOSIS IN TBI

Two prognostic scoring systems for TBI have been developed from the analysis of large patient datasets: the International Mission on Prognosis and Analysis of Clinical trials in Traumatic Brain Injury database (IMPACT) and the Corticosteroid Randomization After Significant Head injury trial data (CRASH) (34, 35). IMPACT and CRASH are based on multiple logistic regression models of variables predicting functional outcome at 6 months (Table 1). Independently predictive variables cluster around clinical signs (e.g., pupil reactivity and motor responses), neuroanatomic descriptors from cranial CT (midline shift and encroachment of basilar cisterns), and selected physiologic and biochemical derangements (Table 1). Both models discriminate outcomes with areas under the receiver operating characteristic curve (AUC) of 0.6–0.8 and have been

calibrated internally and externally (35, 36) (Table 2). The external validity of IMPACT and CRASH models was confirmed in an analysis of five more recent studies totaling 9,036 patients, an analysis which did not demonstrate any meaningful difference in prognostic performance between models (36).

Although extensively validated in large datasets, the CRASH and IMPACT scoring systems do not have the accuracy to be meaningful for decision making at the individual patient level. Prognostic estimates always represent probabilities and not absolute certainties on the actual outcome at any given time. The value of IMPACT and CRASH is more apparent in the design and analysis of clinical trials, where they can be used to stratify patients into a priori prognostic categories, and in clinical audits, where they can be used to adjust for case mix.

Genetic Factors

Research suggests that the APOE ε4 allele, which has been linked to Alzheimer’s disease, may significantly influence outcome

TABLE 2. Validation of the International Mission on Prognosis and Analysis of Clinical Trials and Adapted Corticoid Randomization After Significant Head Injury Models

Reference/ Country	Study Design	Patients	Definition of Poor Outcome	Prediction of Poor Outcome
Steyerberg et al (34)/ The Netherlands	Data from eight randomized controlled trials and three observational studies	8,509 patients with moderate and severe TBI	GOS 1–3 and death at 6 mo	AUC 0.66–0.84 for core model AUC increased by approximately 0.05 with extended model External validation on 6681 patients with AUC 0.78–0.80
Perel et al (35)/United Kingdom	Prospective multicenter randomized trial	10,008 patients	GOS 1–3 at 6 mo and death at 14 d	Basic model: AUC 0.86 for death and 0.81 for GOS 1–3 Extended model: AUC 0.88 for death and 0.83 for GOS 1–3 External validation on 8509 patients with 0.77 for both models
Roozenbeek et al (36)/ The Netherlands	Data from three randomized controlled trials and two observational studies	9,036 patients from five different large datasets	GOS 1–3 and death at 6 mo	External validation of IMPACT against all five datasets showed good discrimination for death (AUC 0.65–0.83) and GOS 1–3 (AUC 0.66–0.76) External validation of CRASH against all five datasets showed good discrimination for death (AUC 0.66–0.85) and GOS 1–3 (AUC 0.68–0.78)

IMPACT = International Mission on Prognosis and Analysis of Clinical Trials; CRASH = Corticoid Randomization After Significant Head injury; TBI = traumatic brain injury; AUC = area under the curve; GOS = Glasgow Outcome Scale.

following TBI (37–50). Distinct neuropathological features, including amyloid deposits, have been found in patients with the APOE ϵ 4 allele who die of TBI (51). Prospective evaluations demonstrate that TBI patients with the APOE ϵ 4 allele are twice as likely to have an unfavorable outcome compared with those without the allele (37, 38). TBI patients with the ϵ 4 allele have larger parenchymal hematomas (41) and are more likely to remain in prolonged coma (39) and to have posttraumatic seizures (42) and adverse rehabilitation outcome (39, 43). Available data indicate that APOE ϵ 4 is a robust predictor of impaired neuropsychological function and of dementia following TBI (44, 45, 47, 48). Other genes have been linked with outcome after TBI, such as APOE promoter (52), genes encoding for the catechol-o-methyltransferase (53), the dopamine D2 receptor (DRD2) (54), interleukin genes (55), the p53 gene that regulates cell cycle (56, 57), PARP-1 (58), and CACNA1A genes (59). While these observations have generated important hypotheses regarding the biology of TBI and its outcome, broader efforts are needed to capture genome-wide expression patterns in large populations of TBI patients.

Clinical Assessment

Clinical estimation of injury severity has long been the cornerstone outcome prediction after TBI (60) and typically centers on an evaluation of level of consciousness and of brainstem

responses (61). The Glasgow Coma Scale (GCS) has been widely adopted as a simple method to numerically express the clinically observed features of consciousness (62). The GCS has several limitations: it does not directly assess brainstem responses, and the value of the verbal and eye subscores is diminished in intubated, aphasic, or aphonic patients and in those who have facial or ocular injury impeding eye evaluation. On the other hand, the GCS motor subscore has robust prognostic value in both CRASH and IMPACT models. Extensor or absent motor responses are associated with poor 6-month functional outcome (35, 63). The more recent Full Outline of UnResponsiveness (FOUR) attempts to address some of the limitations of the GCS by removing any verbal assessment and integrating brainstem responses and breathing patterns (64) (Fig. 2). In a recent prospective evaluation of 51 patients with TBI, the AUC of the FOUR score in predicting poor functional outcome (dead, vegetative, or severely disabled) at 6 months was 0.85, equivalent to the AUC of 0.83 observed with the GCS (65). The FOUR scale allows the explicit testing of eye movements or blinking that facilitates the detection of locked-in state or the transition from VS to MCS. However, the usefulness of a scale that excludes verbal assessment could be debated especially in the TBI population, the majority of whom have moderate or mild injury in which a critical distinguishing feature is verbal performance. Furthermore, respiratory patterns

FOUR Scale	GCS
Eye opening 4 = eyelids open or opened, tracking, or blinking to command 3 = eyelids open but not tracking 2 = eyelids closed but open to loud voice 1 = eyelids closed but open to pain 0 = eyelids remain closed with pain	Eye opening 4 = eyes open spontaneously 3 = eye opening to verbal command 2 = eye opening to pain 1 = no eye opening
Motor response 4 = thumbs-up, fist, or peace sign 3 = localizing to pain 2 = flexion response to pain 1 = extension response to pain 0 = no response to pain or generalized myoclonus status	Motor response 6 = obeys commands 5 = localizing pain 4 = withdrawal from pain 3 = flexion response to pain 2 = extension response to pain 1 = no motor response
Brainstem reflexes 4 = pupil and corneal reflexes present 3 = one pupil wide and fixed 2 = pupil or corneal reflexes absent 1 = pupil and corneal reflexes absent 0 = absent pupil, corneal, and cough reflex	Verbal response 5 = oriented 4 = confused 3 = inappropriate words 2 = incomprehensible sounds 1 = no verbal response
Respiration 4 = not intubated, regular breathing pattern 3 = not intubated, Cheyne-Stokes breathing pattern 2 = not intubated, irregular breathing 1 = breathes above ventilator rate 0 = breathes at ventilator rate or apnea	

Figure 2. Comparison of the Full Outline of UnResponsiveness (FOUR) scale with the Glasgow Coma Scale (GCS). Adapted from Wijdicks et al (64).

can be difficult to interpret in patients undergoing mechanical ventilation. These characteristics suggest that the clinical utility of the FOUR scale may be more relevant in the postacute phase of SBI.

Neuroimaging

Several features readily identified on cranial CT scan have major prognostic significance. These include shift of midline structures, encroachment of the basal cisterns, cerebral infarction, subarachnoid hemorrhage, intraventricular hemorrhage, diffuse injury, and extra-axial hematomas (66–69). Widespread availability and speed of image processing have established CT as a cornerstone in the evaluation and outcome prediction of patients with TBI. However, CT lacks the resolution to correctly identify or characterize smaller white matter lesions as they might occur in the setting of diffuse axonal injury (70).

The significance of brain MRI as a prognostic tool in the acute setting of TBI is being actively investigated (71). MRI sequences such as diffusion weighted imaging and susceptibility weighted imaging are potentially more sensitive to diffuse axonal injury and have been shown to increase the accuracy of outcome prediction (72–74). Recent work indicates that diffusion tensor imaging (DTI) is highly sensitive to traumatic white matter damage (75) and is predictive of long-term functional outcome (76, 77). In a prospective evaluation of 105 patients who were comatose 1–3 weeks after TBI, a prognostic score that integrated DTI measurements

of white matter tracts with the IMPACT score was able to classify 1-year functional outcomes with greater accuracy than the IMPACT score alone (78). Using early proton magnetic resonance spectroscopy, it is possible to identify axonal loss (reduced N-acetyl aspartate-to-creatine ratio) and increased myelin turnover (increased choline-to-creatine ratio) in the tissue that appears normal using conventional morphological MRI sequences (79), changes that have been linked to worse long-term outcome (76, 79, 80).

Brain Physiology and Metabolism

Sustained elevations of intracranial pressure (ICP) to greater than 20 mm Hg or decreases in cerebral perfusion pressure (CPP) to less than 50–60 mm Hg have been linked with cerebral infarction, herniation, and death following TBI (69, 81).

Studies indicate that changes in ICP cannot be predicted in a reliable or timely manner with clinical assessment or imaging (82–85); however, the efficacy of ICP-guided management has been challenged (86–88). Brain tissue monitoring indicates that even when ICP and CPP are within normal limits, significant reductions in brain tissue oxygen pressure (P_{btO_2}) may occur, and brain tissue hypoxia has been associated with worse outcome (89, 90). Consistent with these observations, studies suggest beneficial effects of clinical algorithms that target normalization of P_{btO_2} levels (91–93). Similarly, the use of cerebral microdialysis probes demonstrates neurometabolites in the brain interstitium whose concentrations have been associated with outcome following neurologic trauma (94, 95). In the largest of these studies, a cohort of 223 patients with severe TBI, low brain extracellular glucose and elevated lactate to pyruvate (L/P) were independently predictive of mortality at 6 months (96). Microdialytic assessment may identify irreversible neuronal loss, as indicated in a study that found an association between elevations in L/P in the acute setting and subsequent volume loss in the frontal lobes (97). Preliminary microdialysis studies in patients with severe TBI suggest that brain extracellular concentrations of amyloid fragments, tau protein, and neurofilament heavy chain protein may also have prognostic significance (98–100).

Electrophysiology

The use of continuous electroencephalography (cEEG) suggests that seizure activity and status epilepticus, often clinically

undetected, occur in a significant number of patients with severe TBI, and such perturbations have been linked to outcome (101, 102). In a study of 94 patients with moderate to severe TBI who underwent cEEG monitoring, seizures occurred in 21, and mortality was 100% in patients with status epilepticus (103). Seizures following TBI are associated with increased cerebral metabolic rate, cerebral blood flow and volume, and increasing ICP in susceptible patients (104). It has been suggested that seizures cause injury, compounding trauma-induced damage: hippocampal atrophy was more pronounced in TBI patients with seizures than in those without seizures, and atrophy was most apparent on the hippocampus ipsilateral to the seizure focus (105). Additional research is needed to determine if early and intensive suppression of seizure activity or periodic discharges may improve outcome after TBI; current guidelines recommend 7 days of seizure prophylaxis following moderate to severe TBI (106).

The value of somatosensory evoked potentials (SSEPs) in predicting outcome of severe TBI was evaluated in a meta-analysis of 44 studies published from 1976 to 2000 (107). The bilateral absence of cortical signals on SSEP had a positive predictive value of 98.7% in predicting adverse outcome 2 months to 3 years after head injury; however, the accuracy of prediction is significantly lower when patients have focal lesions, subdural fluid collections, and recent decompressive surgery (107). In a more recent report, bilaterally absent cortical SSEP responses assessed on the third day after severe TBI were correlated with functional outcomes at 1 year; furthermore, SSEP helped predict performance on tests of information-processing speed, working memory, and attention 1 year after injury (108).

Serum Biomarkers

Serum levels of S100 beta protein and neuron-specific enolase (NSE), markers of glial and neuronal damage, respectively, have been extensively studied in patients with TBI. In some studies, these markers correlated with findings on neuroimaging but were not independently predictive of specific outcomes (109–111). Other investigations have indicated that S100 beta and NSE have value not only in determining the severity of TBI but also in classifying their outcomes (112–119). Glial fibrillary acidic protein (GFAP) is specific to glial cells in the central nervous system, and increased levels are detectable in the serum of patients with acquired brain injury. Studies suggest that serum GFAP levels correlate with clinical and neuroradiologic injury severity and are significantly higher in patients who die or have poor functional outcome (116, 119–121). The predictive value of other serum biomarkers, in particular tau protein, is still under investigation (122, 123).

Systemic Biochemical Variables

Several studies have investigated associations of routinely measured laboratory parameters and outcome in patients with TBI. Among them, coagulation disturbances, low platelet counts, as well as anemia, hypoalbuminemia, elevated serum creatinine, and high serum glucose concentrations have been

associated with adverse outcomes in both univariate and multivariate models (63, 124–129). Given the adverse association of high glucose concentrations with outcome, recent randomized trials have evaluated intensive insulin therapy to maintain normoglycemia in TBI patients; however, a distinct outcome benefit was not demonstrated (130–132). Other studies using cerebral microdialysis have found that intensive insulin therapy may result in unfavorable trends in brain extracellular glucose and L/P ratios (133–135). In a randomized crossover trial of intensive (80–110 mg/dL) vs. less intensive (120–150 mg/dL) glycemic control in 13 patients with severe TBI, critical reductions in brain interstitial glucose and elevations of L/P ratio were more frequent in patients allocated to intensive glycemic control and were associated with increased [¹⁸F]-deoxy-D-glucose uptake on PET (136). Given data linking poor glycemic control and adverse head injury outcome (63, 129), additional studies are needed to evaluate whether increased brain glucose delivery via carefully titrated moderate hyperglycemia represents a viable strategy in severe TBI.

Therapeutic Intervention

Evidence suggests that prognosis of TBI may be favorably influenced by rigorous supportive management and enhancements in the process of care. Aggressive measures to normalize oxygenation, volume status, and blood pressure in the early stages of injury have been linked to improved outcome (137–139), as those measures are used in high-volume trauma centers and protocol-driven medical management (140, 141). However, studies favoring these interventions are associative in nature, and direct evidence of their causal effect on the biology and natural history of TBI is lacking.

Although preclinical data on therapeutic hypothermia as a neuroprotectant have been encouraging, a beneficial effect on post-TBI neurological and functional outcomes has not been confirmed in human randomized trials (142, 143). Hypothermia may have greater value in the management of intracranial hypertension. In a systematic review of hypothermia trials in TBI, four of five studies targeting ICP elevation reported a decrease in mortality or in the percentage of patients having a poor recovery, whereas none of the three studies in which hypothermia was assessed as a neuroprotectant showed any benefit (144).

PROGNOSIS AFTER CA

In contrast to the IMPACT and CRASH models for TBI, predictive scoring systems have not been validated in patients with hypoxic-ischemic encephalopathy following CA. Prognosis is preponderantly based on associations derived from physical examination, electrophysiological tests, neuroimaging, and biomarkers. In a statement published in 2006, the American Academy of Neurology expert panel generated recommendations regarding outcome prediction in comatose survivors of CA (145). Based on a review of the literature, the panel concluded that unfavorable outcome can be predicted 72 hours post-CA on the basis of absent pupillary or corneal responses,

TABLE 3. Predictive Value of Clinical Examination in Hypoxic-Ischemic Brain Injury

Reference/ Country	Study Design	Patients or Studies	Time of Examination	Assessment	Definition of Poor Outcome	Prediction of Poor Outcome
Pupillary or corneal reactivity						
Zandbergen et al (199)/The Netherlands	Systematic review	33 studies	In the first 3 d after CPR	Absence of pupillary reactivity	Death or vegetative state	False-positive rate 9%
Booth et al (200)/ Canada	Systematic review	11 studies	In the first 24 hr after CPR	Absence of pupillary reactivity	CPC 3–5	Likelihood ratio 10.2; 95% CI 1.8–48.6
Zandbergen et al (173)/The Netherlands	Prospective study	407 patients from 32 ICUs	In 12, 24, and 72 hr after CPR	Absence of pupillary or corneal reactivity	Death or vegetative state at 1 mo	False-positive rate 4% at 12 hr, 2% at 24 hr, 0% at 72 hr
Wijdicks et al (145)/United States	Systematic review	12 studies	In the first 3 d after CPR	Absence of pupillary reactivity	Death or coma after 1 mo or death, coma, or severe disability requiring nursing care after 6 mo	False-positive rate 0%
Al Thenayan et al (201)/ Canada	Retrospective study	37 patients	At day 3 after CPR and hypothermia	Absence of pupillary or corneal reactivity	No recovery of awareness during hospital stay	False-positive rate 0%
Rossetti et al (202)/ Switzerland	Prospective study	111 patients	In the first 3 d after CPR and hypothermia	Absence of brainstem reflexes	Death during hospital stay	False-positive rate 4%
Fugate et al (187)/United States	Prospective study	103 patients	At day 3 after CPR and hypothermia	Absence of pupillary or corneal reactivity	Death during hospital stay	False-positive rate 0%
Bouwes et al (189)/The Netherlands	Prospective multicenter study (ten centers)	196 patients with pupillary tests; 130 with corneal tests	In the first 3 d after CPR	Absence of pupillary or corneal reactivity	GOS 1–3 at 6 mo	False-positive rate 1% for pupils; 4% for cornea
Rossetti et al (190)/ Switzerland	Prospective study	61 patients	After CPR and hypothermia	Absence of brainstem reflexes	CPC 3–5 at 3 mo	False-positive rate 11%
Motor responses						
Zandbergen et al (199)/ Netherlands	Systematic review	33 studies	In the first 3 d after CPR	Absence of motor responses	Death or vegetative state	False-positive rate 7%
Booth et al (200)/ Canada	Systematic review	1,914 patients from 11 studies	In the first 3 d after CPR	Absence of motor responses	CPC 3–5	Likelihood ratio 9.2; 95% CI 2.1–49.4
Zandbergen et al (173)/The Netherlands	Prospective study	407 patients from 32 ICUs	At 12, 48, and 72 hr after CPR	Absence of motor responses	Death or vegetative state at 1 mo	False-positive rate 9% at 12 hr, 6% at 24 hr, 5% at 72 hr

(Continued)

TABLE 3. (Continued). Predictive Value of Clinical Examination in Hypoxic-Ischemic Brain Injury

Reference/ Country	Study Design	Patients or Studies	Time of Examination	Assessment	Definition of Poor Outcome	Prediction of Poor Outcome
Wijdicks et al (145)/United States	Systematic review	12 studies	In the first 3 d after CPR	GCS motor score 1–2	Death or coma after 1 mo or death, coma, or severe disability requiring nursing care after 6 mo	False-positive rate 0%
Al Thenayan et al (201)/ Canada	Retrospective study	37 patients	At day 3 after CPR and hypothermia	Motor response no better than extension	No recovery of awareness during hospital stay	False-positive rate 14%
Rossetti et al (202)/ Switzerland	Prospective study	111 patients	In the first 3 d after CPR and hypothermia	Absence of motor responses	Death during hospital stay	24% with false-positive mortality predictions during hypothermia
Fugate et al (187)/United States	Prospective study	70 patients (of 103)	At day 3 after CPR and hypothermia	Motor response no better than extension	Death during hospital stay	False-positive rate 3%
Bouwes et al (189)/The Netherlands	Prospective multicenter study (ten centers)	284 patients (of 391)	At day 3 after CPR and hypothermia	Motor response no better than extension	GOS 1–3 at 6 mo	False-positive rate 10%
Rossetti et al (190)/ Switzerland	Prospective study	61 patients	After CPR and hypothermia	Extension or no motor response	CPC 3–5 at 3 mo	False-positive rate 7%
Myoclonus						
Wijdicks et al (203)/United States	Prospective study	107 patients	After CPR	Presence of myoclonus	Death during hospital stay	False-positive rate 0%
Zandbergen et al (173)/The Netherlands	Prospective study	407 patients from 32 ICUs	In first 3 d after CPR	Presence of myoclonus	Death or vegetative state at 1 mo	False-positive rate 0%
Wijdicks et al (145)/United States	Systematic review	12 studies	In the first 3 d after CPR and hypothermia	Presence of myoclonus	Death or coma after 1 mo or death, coma, or severe disability requiring nursing care after 6 mo	False-positive rate 0%
Al Thenayan et al (201)/ Canada	Retrospective study	37 patients	At day 3 after CPR and hypothermia	Presence of myoclonus	No recovery of awareness during hospital stay	False-positive rate 0%
Rossetti et al (202)/ Switzerland	Prospective study	111 patients	In the first 3 d after CPR and hypothermia	Presence of myoclonus	Death during hospital stay	7% with false- positive mortality predictions during hypothermia
Fugate et al (187)/United States	Prospective study	103 patients	At day 3 after CPR and hypothermia	Presence of myoclonus	Death during hospital stay	False-positive rate 0%
Rossetti et al (190)/ Switzerland	Prospective study	61 patients	After CPR and hypothermia	Presence of myoclonus	CPC 3–5 at 3 mo	False-positive rate 7%

CPR = cardiopulmonary resuscitation; CPC = Cerebral Performance Categories Scale; GOS = Glasgow Outcome Scale; CI = confidence interval; GCS = Glasgow Coma Scale.

TABLE 4. Predictive Value of Neuron-Specific Enolase for Poor Outcome in Hypoxic-Ischemic Brain Injury

Reference/ Country	Study Design	Patients or Studies	Time of Examination	Assessment	Definition of Poor Outcome	Prediction of Poor Outcome
NSE						
Zandbergen et al (204)/The Netherlands	Systematic review	802 patients from 28 studies	After CPR	NSE in cerebrospinal fluid > 33 ng/ mL	Death or persistent vegetative state	False-positive rate 0%
Meynaar et al (185)/ Netherlands	Prospective study	110 patients	At 6 hr after CPR	Serum NSE > 25 ng/mL	Death or no recovery from coma until discharge	False-positive rate 0%
Zingler et al (205)/ Germany	Prospective study	27 patients	At days 1, 2, 3, and 7 after CPR	Serum NSE at different cutoffs	CPC 3–5 at 3 mo	False-positive rate 0% at different cutoffs on all days
Tiainen et al (186)/ Finland	Prospective study	70 patients	At 24 and 36 hr after CPR with or without hypothermia	Serum NSE at different cutoffs	CPC 3–5 at 6 mo	At all time points and all cutoffs false- positive rate 0% in patients without, 4% in patients with hypothermia. Lower levels with hypothermia
Zandbergen et al (173)/The Netherlands	Prospective study	407 patients from 32 ICUs	At 24, 48, and 72 hr after CPR	Serum NSE > 33 ng/mL	Death or vegetative state at 1 mo	False-positive rate 0% at all time points
Wijdicks et al (145)/ United States	Systematic review	12 studies	At day 3 after CPR	Serum NSE at different cutoffs	Death or coma after 1 mo or death, coma, or severe disability requiring nursing care after 6 mo	False-positive rate 0% at different cutoffs
Fugate et al (187)/ United States	Prospective study	74 patients (of 103)	At day 3 after CPR and hypothermia	Serum NSE > 33 ng/mL	Death during hospital stay	False-positive rate 29%
Steffen et al (188)/ Germany	Prospective study	230 patients	At 72 h after CPR with and without hypothermia	Serum NSE at different cutoffs	CPC 3–5 at discharge	ROC analysis: higher cutoff with a false- positive rate 0% in hypothermia (78.9 ng/mL) compared to no hypothermia (26.9 ng/mL)
Cronberg et al (206)/ Sweden	Prospective study	34 patients (of 111)	At 48 hr after CPR and hypothermia	Serum NSE > 33 ng/mL	CPC 3–5 at 6 mo	False-positive rate 0%
Bouwes et al (189)/The Netherlands	Prospective multicenter study (ten centers)	391 patients	At 12, 36, and 48 hr after CPR and hypothermia	Serum NSE > 33 ng/mL	GOS 1–3 at 6 mo	False-positive rate 10% at 12 hr, 9% at 36 hr, and 7% at 48 hr
Rossetti et al (190)/ Switzerland	Prospective study	61 patients	At 24 and 48 hr after CPR and hypothermia	Serum NSE > 33 ng/mL	CPC 3–5 at 3 mo	False-positive rate 4%

CPR = cardiopulmonary resuscitation; NSE = neuron-specific enolase; ROC = receiver operating characteristic; CPC = Cerebral Performance Categories Scale; GOS = Glasgow Outcome Scale.

TABLE 5. Predictive Value of Electrophysiological Assessment in Hypoxic-Ischemic Brain Injury

Reference/ Country	Study Design	Patients or Studies	Time of Examination	Assessment	Definition of Poor Outcome	Prediction of Poor Outcome
Somatosensory evoked potentials						
Chen et al (167)/ Canada	Prospective case series	34 patients	In the first 3 d after CPR	Bilateral absence of N20	No recovery, coma until death or persistent vegetative state	False-positive rate 0%
Pohlmann-Eden et al (207)/ Germany	Prospective study	42 patients	After 36 hr after CPR	Bilateral absence of N20	GOS 3–5 at 3 mo	False-positive rate 3%
Zandbergen et al (199)/The Netherlands	Systematic review	33 studies	In the first week after CPR	Bilateral absence of N20	Death or vegetative state	False-positive rate 0%
Meynaar et al (185)/The Netherlands	Prospective study	59 patients (of 110)	At ≥ 48 hr after CPR	Bilateral absence of N20	Death or no recovery from coma until discharge	False-positive rate 0%
Zandbergen et al (173)/The Netherlands	Prospective study	407 patients from 32 ICUs	At 24, 48, and 72 hr after CPR	Bilateral absence of N20	Death or vegetative state at 1 mo	False-positive rate 0% for all time points
Wijdicks et al (145)/United States	Systematic review	12 studies	At day 3 after CPR	Bilateral absence of N20	Death or coma after 1 mo or death, coma, or severe disability requiring nursing care after 6 mo	False-positive rate 0%
Bouwes et al (208)/The Netherlands	Prospective multicenter study (two centers)	77 patients	After CPR and during hypothermia	Bilateral absence of N20	GOS 1–2 at 1 mo	False-positive rate 0% during hypothermia
Rossetti et al (202)/ Switzerland	Prospective study	111 patients	In the first 3 d after CPR and hypothermia	Bilateral absence of N20	Death during hospital stay	False-positive rate 0%
Fugate et al (187)/United States	Prospective study	14 patients (of 103)	After CPR and during hypothermia	Bilateral absence of N20	Death during hospital stay	False-positive rate 0% during hypothermia
Leithner et al (209)/ Germany	Retrospective study	112 patients (of 185)	After CPR and during hypothermia	Bilateral absence of N20	CPC 3–5 at discharge	1/36 patients with absent early N20 had good outcome during hypothermia
Bouwes et al (189)/The Netherlands	Prospective multicenter study (ten centers)	391 patients	After CPR and during and after hypothermia	Bilateral absence of N20	GOS 1–3 at 6 mo	False-positive rate 3% during hypothermia; 0% after hypothermia
Rossetti et al (190)/ Switzerland	Prospective study	61 patients	After CPR and hypothermia	Bilateral absence of N20	CPC 3–5 at 3 mo	False-positive rate 0%

(Continued)

TABLE 5. (Continued). Predictive Value of Electrophysiological Assessment in Hypoxic-Ischemic Brain Injury

Reference/ Country	Study Design	Patients or Studies	Time of Examination	Assessment	Definition of Poor Outcome	Prediction of Poor Outcome
EEG patterns Chen et al (167)/ Canada	Prospective case series	34 patients	In the first 3 d after CPR	Suppression, unreactive alpha/theta, epileptiform discharges, burst- suppression.	No recovery, coma until death or persistent vegetative state	Two patients had good outcome
Zandbergen et al (199)/The Netherlands	Systematic review	33 studies	After CPR	EEG burst- suppression or isoelectric trace	Death or vegetative state	False-positive rate 9%
Zandbergen et al (173)/The Netherlands	Prospective study	407 patients from 32 ICUs	After CPR	EEG burst- suppression or isoelectric curve; postanoxic status epilepticus	Death or vegetative state at 1 mo	False-positive rate 0% for EEG burst- suppression or isoelectric curve; false- positive rate for 7% postanoxic status epilepticus
Wijdicks et al (145)/United States	Systematic review	12 studies	After CPR	EEG burst- suppression or generalized epileptiform discharges	Death or coma after 1 mo or death, coma, or severe disability requiring nursing care after 6 mo	False-positive rate 3%
Rossetti et al (210)/ Switzerland	Retrospective study	107 patients (of 166)	After CPR with and without hypothermia	Postanoxic status epilepticus	Death during hospital stay	False-positive rate 8%
Rossetti et al (171)/ Switzerland	Prospective study	34 patients	After CPR and during hypothermia	cEEG burst- suppression, seizures or epileptiform discharges	Death during hospital stay	False-positive rate 0% during hypothermia
Rossetti et al (202)/ Switzerland	Prospective study	111 patients	In the first 3 d after CPR and hypothermia	EEG epileptiform activity	Death during hospital stay	False-positive rate 9%
Fugate et al (187)/United States	Prospective study	21 patients (of 103)	At day 3 after CPR and hypothermia	EEG burst- suppression or isoelectric curve. Status epilepticus or no background reactivity	Death during hospital stay	False-positive rate 0%
Thenayan et al (211)/Saudi Arabia	Retrospective study	29 patients	After CPR with and without hypothermia	EEG with generalized suppression or burst- suppression	No recovery of awareness during hospital stay	False-positive rate 0%
Rossetti et al (190)/ Switzerland	Prospective study	61 patients	After CPR and during hypothermia	Epileptiform transients	CPC 3–5 at 3 mo	False-positive rate 0% during hypothermia

(Continued)

TABLE 5. (Continued). Predictive Value of Electrophysiological Assessment in Hypoxic-Ischemic Brain Injury

Reference/ Country	Study Design	Patients or Studies	Time of Examination	Assessment	Definition of Poor Outcome	Prediction of Poor Outcome
EEG background reactivity						
Rossetti et al (171)/ Switzerland	Prospective study	34 patients	After CPR and during hypothermia	Absence of cEEG background reactivity	Death during hospital stay	False-positive rate 0% during hypothermia
Rossetti et al (202)/ Switzerland	Prospective study	111 patients	In the first 3 d after CPR and hypothermia	Absence of EEG background reactivity	Death during hospital stay	False-positive rate 7%
Thenayan et al (211)/ Saudi Arabia	Retrospective study	29 patients	After CPR and with or without hypothermia	Preserved EEG background reactivity	No recovery of awareness during hospital stay	10/11 patients with reactivity regained awareness
Rossetti et al (190)/ Switzerland	Prospective study	61 patients	After CPR and during hypothermia	Absence of EEG background reactivity	CPC 3–5 at 3 mo	False-positive rate 0% during and after hypothermia
Howard et al (212)/United Kingdom	Retrospective study	39 patients	At a mean of 5 d after CPR	Absence of EEG background reactivity or periodic generalized phenomenon	Death, profound cognitive impairment, including persistent vegetative state, minimally area states, or severe physical impairment at discharge	Significant association with poor outcome (false-positive rate not provided)

CPR = cardiopulmonary resuscitation; EEG = electroencephalography; cEEG = continuous electroencephalography; GOS = Glasgow Outcome Scale; CPC = Cerebral Performance Categories Scale.

or absent or extensor motor response (**Table 3**). The panel also pointed to myoclonus status epilepticus, elevated serum NSE (**Table 4**), and absent cortical response of SSEPs (**Table 5**) as additional predictive elements. These recommendations were exclusively based on studies of patients who were not treated with therapeutic hypothermia.

Prognostication following CA has been considered in terms of prearrest factors, intra-arrest factors, and postresuscitation factors (6, 146). Prearrest factors include age, race, Acute Physiology and Chronic Health Evaluation II and III scores, diabetes mellitus, sepsis, metastatic cancer, renal failure, homebound lifestyle, and stroke; these factors have been linked with outcome but are not reliable prognostic variables (6). Intra-arrest factors include the duration of time between circulatory collapse and the start of cardiopulmonary resuscitation (CPR), duration of CPR to return of spontaneous circulation, asystole, noncardiac causes of arrest, nonadherence to established CPR guidelines, and maximum end-tidal carbon dioxide (ET_{CO₂}) of less than 10 mm Hg—all linked to unfavorable outcome.

Postresuscitation factors are discussed below.

There is a need for large-scale prospective studies designed to validate the individual and integrated significance of clinical, biochemical, electrophysiological, and imaging variables in predicting CA outcome. Meanwhile, existing recommendations regarding prognostication should be applied with caution in CA patients treated with hypothermia, as the accuracy of tests during and after hypothermia has been seriously challenged. An approach for prognostication in CA patients treated with hypothermia is shown in **Figure 3**.

Clinical Assessment

As in patients with TBI, outcome prediction in hypoxic-ischemic encephalopathy is heavily reliant on clinical assessment. In a landmark early study conducted in patients with coma from a range of nontraumatic etiologies, unfavorable 1-year outcome could be predicted on the basis of a limited number of clinical signs, including the inability to follow commands, the lack of visual fixation or pursuit,




Figure 3. Proposed algorithm for prognostication in hypoxic-ischemic brain injury. SSEPs = somatosensory evoked potentials; FPR = false-positive rate; EEG = electroencephalography; ADC = apparent diffusion coefficient; cEEG = continuous electroencephalography; NSE = neuron-specific enolase; ROSC = return of spontaneous circulation. *Definitions of unfavorable outcome are presented in Tables 3–6.

and the lack of oculocephalic or oculovestibular responses (147). These results and additional analysis in a subset of patients with hypoxic-ischemic encephalopathy (148) are the evidentiary basis for a widely cited practice guideline from the American Academy of Neurology (145). In a more recent cohort of 500 patients, the pupillary and, to a lesser degree, oculocephalic responses were the clinical findings most predictive of 6-month functional outcome (149). Current prospective evaluations in CA patients who have been treated with hypothermia seriously challenge the prognostic accuracy of selected clinical findings, in particular the association between absent or abnormal motor responses and poor outcome, with false-positive rates of 10% and 24% in two recent reports (150, 151).

Neuroimaging

Neuropathological studies of patients who died following CA indicate neuronal death that is prominent in occipital, frontal, and parietal cortex; the hippocampus; the basal ganglia; the thalamic reticular nucleus; Purkinje cells of the cerebellum; and the spinal cord (152). When evaluated with cranial CT, early hypoxic-ischemic damage may be suggested by a loss of differentiation between gray and white matter (153, 154); however, in many cases, CT is unrevealing. With brain MRI, early reports demonstrated multifocal cortical (155) or white matter changes (156) that were associated with poor outcome following CA (155–157). More recently, the prognostic value

of quantitative diffusion-weighted MRI has been studied in CA survivors (158–160). In a prospective evaluation of 51 patients, one group found that whole-brain apparent diffusion coefficients (ADCs) derived from MRI 49 to 108 hours after CA accurately classified 6-month functional outcome, surpassing the sensitivity of clinical neurological assessment (159). Related studies indicate that unfavorable outcome might be predicted by reduced regional ADC values in the putamen and in occipital, parietal, and temporal cortices (158, 161) (Table 6). In a recent prospective evaluation of 57 CA patients, it was found that a prognostic model based on a quantitative DTI assessment of white matter tracts predicted the 1-year functional outcome with a high degree of accuracy (162). Anecdotal reports in patients with CA

indicate that brain proton magnetic resonance spectroscopy identifies metabolic abnormalities associated with neuronal and glial cell damage; however, the prognostic significance of these findings requires further study (163, 164).

Electrophysiologic Tests

Seizures are detected in up to 40% of comatose survivors of CA and have been linked to adverse outcome (146, 148, 165, 166). Malignant EEG patterns, such as burst-suppression, nonreactive, or flat-line EEG, are powerful indicators of unfavorable prognosis (167–169); however, these findings have reduced specificity in patients treated with hypothermia (150, 151), a limitation that may be overcome with the help of quantitative EEG methods (170). Recently, the presence of background EEG reactivity in response to a noxious stimulus was identified as an important new predictor of favorable outcome (151, 171) (Table 5).

Available studies indicate that short-latency SSEPs have robust prognostic value in patients after CA; the bilateral absence of the cortical N20 potential has a specificity of close to 100% for the prediction of poor outcome (145, 146, 167, 172–174) (Table 5). While absent or abnormal N20 signals are validated predictors of poor outcome, it has been hypothesized that long-latency potentials (e.g., P300 and mismatch negativity [MMN]), which evaluate the functional integrity of cortico-cortical and thalamocortical circuits, might be best suited to predict coma emergence and higher order cognitive function

TABLE 6. Predictive Values of Neuroimaging for Poor Outcome in Hypoxic-Ischemic Brain Injury

Reference/ Country	Study Design	Patients or Studies	Time of Examination	Assessment	Definition of Poor Outcome	Prediction of Poor Outcome
Wijdicks et al (155)/ United States	Prospective study	27 patients	In the first 15 d after CPR	Brain MRI with diffuse signal abnormalities in the cortex and subcortical areas or effacement of the sulci	Death during hospital stay	All eight patients with these MRI findings died; one of two patients who survived had subcortical signs of ischemia
Wu et al (158)/ United States	Retrospective study	80 patients	In first 7 d after CPR and hypothermia	Brain MRI with lower whole brain and regional median ADC	mRS 3–6 at 6 mo	Patients with mRS > 3 had significant lower median whole brain and regional ADC
Topcuoglu et al (213)/ Turkey	Prospective study	22 patients	At a median 4.1 d (good outcome) and 9.8 d (poor outcome) after CPR	Brain MRI with DWI and FLAIR with lesion pattern of multilobar, or diffuse, cortical involvement	CPC 4–5 at discharge	False-positive rate 0%
Wijman et al (159)/ United States	Prospective study	40 patients (of 83)	In the first 7 d after CPR and hypothermia	Brain MRI with ADC < 650 × 10 ⁶ mm ² /s	Death during hospital stay	False-positive rate 0%
Fugate et al (187)/ United States	Prospective study	53 patients (of 103)	At a median 1 d after CPR and hypothermia	Brain CT with global edema	Death during hospital stay	False-positive rate 0%
Choi et al (160)/ Korea	Prospective study	22 patients (of 111)	At 48 hr after CPR and hypothermia	Brain MRI with global ischemia or focal ischemia with total lesion volume > 20 mL	CPC 3–5 at 6 mo	False-positive rate 0%
Cronberg et al (206)/ Sweden	Prospective study	39 patients	In the first 5 d after CPR and hypothermia	Brain MRI with cortical and/or deep gray nuclei lesions	GOS 1–3 at 3 mo	False-positive rate 23%
Howard et al (212)/ United Kingdom	Retrospective study	39 patients	Between 1 and 150 d after CPR	Brain MRI with extensive changes in cortex and the deep gray matter on DWI and T2-weighted imaging	Death, profound cognitive impairment including persistent vegetative state, minimally area states or severe physical impairment at discharge	False-positive rate 0%

CPR = cardiopulmonary resuscitation; DWI = diffusion weighted imaging; ADC = apparent diffusion coefficient; FLAIR = fluid-attenuated inversion recovery; mRS = modified Ranking Scale; CPC = Cerebral Performance Categories Scale; GOS = Glasgow Outcome Scale.

(175, 176). The MMN reflects cortical discrimination of an “oddball” stimulus within a series of identical sounds (177). In a seminal report on 62 comatose CA survivors, all patients in whom MMN was present recovered consciousness (178). In a prospective nonrandomized comparison, the amplitude of the P300 response was significantly higher in CA patients who received therapeutic hypothermia than in those who were normothermic (179).

Serum Biomarkers

Serum and cerebrospinal fluid markers, NSE and S100 beta in particular, have been evaluated in patients with hypoxic-ischemic encephalopathy (180–184). Older studies indicated that elevations in serum NSE were robustly associated with poor outcome (146, 185, 186). More recent work indicates that the accuracy of serum NSE elevation in patients who received therapeutic hypothermia is significantly reduced, with false-positive rates of 4% to 29% (Table 4) (187–190). The prognostic significance of other serum biomarkers is less well studied. Preliminary data suggest that serum GFAP accurately discriminates prognostic categories following CA (191) and that serum procalcitonin may classify outcome with greater accuracy than GFAP (192). In a very recent study, serum levels of tau protein were highly predictive of 6-month post-CA functional outcome (193).

Therapeutic Interventions

Lowering of body temperature to 32°–34°C during the first hours after CA reduces neurologic injury by concurrently disrupting a range of pathological cellular events (194, 195). Two randomized trials published in 2002 demonstrated that therapeutic hypothermia implemented early after CA reduces mortality and significantly improves functional outcome (4, 5). In the Hypothermia After Cardiac Arrest Study, 55% of the cooled patients had favorable functional outcomes at 6 months compared with 39% of normothermic control patients (5); these results were confirmed in a contemporary study from Australia (4). Based on preliminary findings, it was postulated that a shorter delay to reach target temperature would lead to improved functional outcome; however, this was not verified in two recent randomized trials (196, 197).

Therapeutic hypothermia, possibly the first targeted intervention in SBI with an unequivocal outcome-modifying effect, has not surprisingly challenged older principles of CA prognostication. Prior to the era of hypothermia, selected variables were recognized as accurate predictors of poor outcome when observed or measured 72 hours after CA, particularly the absence of brainstem reflexes, the presence of extensor motor responses, and a serum NSE level greater than 33 ng/mL (145). Recent studies in patients treated with hypothermia indicate that these same variables fail to accurately discriminate outcome categories, with unacceptably high false-positive rates for the prediction of poor outcome when used in the same time frame as the older studies (Tables 3–6). Emerging evidence suggests that hypothermia modifies the time course of post-CA neurologic injury and recovery. It follows that the

determination of a CA prognosis, traditionally ascertained at 72 hours, in all likelihood, must be deferred to a later time point. However, there are insufficient data to recommend specifically when prognostic assessment should be made (198). Large prospective studies are needed to build prediction models integrating the neurologic examination and electrophysiological and neuroimaging assessments for outcome after CA.

CONCLUSIONS

In patients with SBI, the goal of accurate and reliable prognosis is to guide treatment strategies, so that they are proportionate with outcome. Models for outcome prediction include data from clinical assessment; physiological status; and laboratory, neuroimaging, and electrophysiological examinations. Existing prognostic systems need to be revised, as treatment strategies, such as therapeutic hypothermia, reduce their predictive value. In addition, determinants of outcome, such as genetic susceptibility and biological heterogeneity, need to be rigorously evaluated. Integration of these factors is likely to increase the accuracy of prognostic models in patients suffering from SBI.

REFERENCES

- Langlois JA, Rutland-Brown W, Wald MM: The epidemiology and impact of traumatic brain injury: A brief overview. *J Head Trauma Rehabil* 2006; 21:375–378
- McNally B, Robb R, Mehta M, et al: Out-of-hospital cardiac arrest surveillance—Cardiac Arrest Registry to Enhance Survival (CARES), United States, October 1, 2005–December 31, 2010. *MMWR Surveill Summ* 2011; 60:1–19
- Merchant RM, Yang L, Becker LB, et al: Incidence of treated cardiac arrest in hospitalized patients in the United States. *Crit Care Med* 2011; 39:2401–2406
- Bernard SA, Gray TW, Buist MD, et al: Treatment of comatose survivors of out-of-hospital cardiac arrest with induced hypothermia. *N Engl J Med* 2002; 346:557–563
- Hypothermia After Cardiac Arrest Study Group. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *N Engl J Med* 2002; 346:549–556
- Neumar RW, Nolan JP, Adrie C, et al: Post-cardiac arrest syndrome: Epidemiology, pathophysiology, treatment, and prognostication. *Circulation* 2008; 118:2452–2483
- Peberdy MA, Callaway CW, Neumar RW, et al: Part 9: Post-cardiac arrest care: 2010 American Heart Association Guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2010; 122(18 Suppl 3):S768–S786
- Sunde K, Pytte M, Jacobsen D, et al: Implementation of a standardised treatment protocol for post resuscitation care after out-of-hospital cardiac arrest. *Resuscitation* 2007; 73:29–39
- Hemphill JC 3rd, White DB: Clinical nihilism in neuroemergencies. *Emerg Med Clin North Am* 2009; 27:27–37
- Stevens RD, Bhardwaj A: Approach to the comatose patient. *Crit Care Med* 2006; 34:31–41
- Medical aspects of the persistent vegetative state (1). The Multi-Society Task Force on PVS. *N Engl J Med* 1994; 330:1499–1508
- Medical aspects of the persistent vegetative state (2). The Multi-Society Task Force on PVS. *N Engl J Med* 1994; 330:1572–1579
- Giaccio JT, Ashwal S, Childs N, et al: The minimally conscious state: Definition and diagnostic criteria. *Neurology* 2002; 58:349–353
- Taylor CM, Aird VH, Tate RL, et al: Sequence of recovery during the course of emergence from the minimally conscious state. *Arch Phys Med Rehabil* 2007; 88:521–525

15. Jennett B, Plum F: Persistent vegetative state after brain damage. A syndrome in search of a name. *Lancet* 1972; 1:734–737
16. Kinney HC, Korein J, Panigrahy A, et al: Neuropathological findings in the brain of Karen Ann Quinlan. The role of the thalamus in the persistent vegetative state. *N Engl J Med* 1994; 330:1469–1475
17. Dougherty JH Jr, Rawlinson DG, Levy DE, et al: Hypoxic-ischemic brain injury and the vegetative state: Clinical and neuropathologic correlation. *Neurology* 1981; 31:991–997
18. Adams JH, Graham DI, Jennett B: The neuropathology of the vegetative state after an acute brain insult. *Brain* 2000; 123 (Pt 7):1327–1338
19. Cruse D, Chennu S, Chatelle C, et al: Bedside detection of awareness in the vegetative state: A cohort study. *Lancet* 2011; 378:2088–2094
20. Monti MM, Vanhaudenhuyse A, Coleman MR, et al: Willful modulation of brain activity in disorders of consciousness. *N Engl J Med* 2010; 362:579–589
21. Coleman MR, Menon DK, Fryer TD, et al: Neurometabolic coupling in the vegetative and minimally conscious states: Preliminary findings. *J Neurol Neurosurg Psychiatr* 2005; 76:432–434
22. Schiff ND, Rodriguez-Moreno D, Kamal A, et al: fMRI reveals large-scale network activation in minimally conscious patients. *Neurology* 2005; 64:514–523
23. Boly M, Faymonville ME, Peigneux P, et al: Auditory processing in severely brain injured patients: Differences between the minimally conscious state and the persistent vegetative state. *Arch Neurol* 2004; 61:233–238
24. Giacino JTK, Kalmar K. The vegetative and minimally conscious states: A comparison of clinical features and functional outcome. *J Head Trauma Rehabil* 1997; 12:36–51
25. Giacino JT, Kalmar K. The vegetative and minimally conscious states: A comparison of clinical features and functional outcome. *J Head Trauma Rehabil* 2005; 12:36–51
26. Ansell BJ, Keenan JE: The Western Neuro Sensory Stimulation Profile: A tool for assessing slow-to-recover head-injured patients. *Arch Phys Med Rehabil* 1989; 70:104–108
27. Shiel A, Horn SA, Wilson BA, et al: The Wessex Head Injury Matrix (WHIM) main scale: A preliminary report on a scale to assess and monitor patient recovery after severe head injury. *Clin Rehabil* 2000; 14:408–416
28. Whyte J, Katz D, Long D, et al: Predictors of outcome in prolonged posttraumatic disorders of consciousness and assessment of medication effects: A multicenter study. *Arch Phys Med Rehabil* 2005; 86:453–462
29. Bhatia KP, Marsden CD: The behavioural and motor consequences of focal lesions of the basal ganglia in man. *Brain* 1994; 117(Pt 4):859–876
30. Fève AP, Fénelon G, Wallays C, et al: Axial motor disturbances after hypoxic lesions of the globus pallidus. *Mov Disord* 1993; 8:321–326
31. Govaerts A, Van Zandijcke M, Dehaene I: Posthypoxic midbrain tremor. *Mov Disord* 1998; 13:359–361
32. Hawker K, Lang AE: Hypoxic-ischemic damage of the basal ganglia. Case reports and a review of the literature. *Mov Disord* 1990; 5:219–224
33. Arciniegas DB: Designing clinical trials to improve neurobehavioral outcome after traumatic brain injury: From bench to bedside. *Crit Care Med* 2009; 37:784–785
34. Steyerberg EW, Mushkudiani N, Perel P, et al: Predicting outcome after traumatic brain injury: Development and international validation of prognostic scores based on admission characteristics. *PLoS Med* 2008; 5:e165
35. Perel P, Arango M, Clayton T, et al: Predicting outcome after traumatic brain injury: Practical prognostic models based on large cohort of international patients. *BMJ* 2008; 336(7641):425–429
36. Roozenbeek B, Lingsma HF, Lecky FE, et al: Prediction of outcome after moderate and severe traumatic brain injury: External validation of the International Mission on Prognosis and Analysis of Clinical Trials (IMPACT) and Corticoid Randomisation After Significant Head Injury (CRASH) prognostic models. *Crit Care Med* 2012; 40:1609–1617
37. Chiang MF, Chang JG, Hu CJ: Association between apolipoprotein E genotype and outcome of traumatic brain injury. *Acta Neurochir (Wien)* 2003; 145:649–653
38. Teasdale GM, Nicoll JA, Murray G, et al: Association of apolipoprotein E polymorphism with outcome after head injury. *Lancet* 1997; 350:1069–1071
39. Friedman G, Froom P, Sazbon L, et al: Apolipoprotein E-epsilon4 genotype predicts a poor outcome in survivors of traumatic brain injury. *Neurology* 1999; 52:244–248
40. Sorbi S, Nacmias B, Piacentini S, et al: ApoE as a prognostic factor for post-traumatic coma. *Nat Med* 1995; 1:852
41. Liaquat I, Dunn LT, Nicoll JA, et al: Effect of apolipoprotein E genotype on hematoma volume after trauma. *J Neurosurg* 2002; 96:90–96
42. Diaz-Arrastia R, Gong Y, Fair S, et al: Increased risk of late posttraumatic seizures associated with inheritance of APOE epsilon4 allele. *Arch Neurol* 2003; 60:818–822
43. Lichtman SW, Seliger G, Tycko B, et al: Apolipoprotein E and functional recovery from brain injury following postacute rehabilitation. *Neurology* 2000; 55:1536–1539
44. Liberman JN, Stewart WF, Wesnes K, et al: Apolipoprotein E epsilon 4 and short-term recovery from predominantly mild brain injury. *Neurology* 2002; 58:1038–1044
45. Crawford FC, Vanderploeg RD, Freeman MJ, et al: APOE genotype influences acquisition and recall following traumatic brain injury. *Neurology* 2002; 58:1115–1118
46. Koponen S, Taiminen T, Kairisto V, et al: APOE-epsilon4 predicts dementia but not other psychiatric disorders after traumatic brain injury. *Neurology* 2004; 63:749–750
47. Ariza M, Pueyo R, Matarin Mdel M, et al: Influence of APOE polymorphism on cognitive and behavioural outcome in moderate and severe traumatic brain injury. *J Neurol Neurosurg Psychiatry* 2006; 77:1191–1193
48. Jordan BD, Relkin NR, Ravdin LD, et al: Apolipoprotein E epsilon4 associated with chronic traumatic brain injury in boxing. *JAMA* 1997; 278:136–140
49. Smith C, Graham DI, Murray LS, et al: Association of APOE e4 and cerebrovascular pathology in traumatic brain injury. *J Neurol Neurosurg Psychiatry* 2006; 77:363–366
50. Teasdale GM, Murray GD, Nicoll JA: The association between APOE epsilon4, age and outcome after head injury: A prospective cohort study. *Brain* 2005; 128(Pt 11):2556–2561
51. Nicoll JA, Roberts GW, Graham DI: Apolipoprotein E epsilon 4 allele is associated with deposition of amyloid beta-protein following head injury. *Nat Med* 1995; 1:135–137
52. Lendon CL, Harris JM, Pritchard AL, et al: Genetic variation of the APOE promoter and outcome after head injury. *Neurology* 2003; 61:683–685
53. Lipsky RH, Sparling MB, Ryan LM, et al: Association of COMT Val-158Met genotype with executive functioning following traumatic brain injury. *J Neuropsychiatry Clin Neurosci* 2005; 17:465–471
54. McAllister TW, Rhodes CH, Flashman LA, et al: Effect of the dopamine D2 receptor T allele on response latency after mild traumatic brain injury. *Am J Psychiatry* 2005; 162:1749–1751
55. Uzan M, Tanriverdi T, Baykara O, et al: Association between interleukin-1 beta (IL-1beta) gene polymorphism and outcome after head injury: An early report. *Acta Neurochir (Wien)* 2005; 147:715–720
56. Fisher DE: The p53 tumor suppressor: Critical regulator of life & death in cancer. *Apoptosis* 2001; 6:7–15
57. Martinez-Lucas P, Moreno-Cuesta J, Garcia-Olmo DC, et al: Relationship between the Arg72Pro polymorphism of p53 and outcome for patients with traumatic brain injury. *Intensive Care Med* 2005; 31:1168–1173
58. Sarnaik AA, Conley YP, Okonkwo DO, et al: Influence of PARP-1 polymorphisms in patients after traumatic brain injury. *J Neurotrauma* 2010; 27:465–471
59. Kors EE, Terwindt GM, Vermeulen FL, et al: Delayed cerebral edema and fatal coma after minor head trauma: Role of the CACNA1A calcium channel subunit gene and relationship with familial hemiplegic migraine. *Ann Neurol* 2001; 49:753–760

60. Lingsma HF, Roozenbeek B, Steyerberg EW, et al: Early prognosis in traumatic brain injury: From prophecies to predictions. *Lancet Neurol* 2010; 9:543–554
61. Jennett B, Teasdale G, Braakman R, et al: Prognosis of patients with severe head injury. *Neurosurgery* 1979; 4:283–289
62. Teasdale G, Jennett B: Assessment of coma and impaired consciousness. A practical scale. *Lancet* 1974; 2:81–84
63. Murray GD, Butcher I, McHugh GS, et al: Multivariable prognostic analysis in traumatic brain injury: Results from the IMPACT study. *J Neurotrauma* 2007; 24:329–337
64. Wijdicks EF, Bamlet WR, Maramattom BV, et al: Validation of a new coma scale: The FOUR score. *Ann Neurol* 2005; 58:585–593
65. Sadaka F, Patel D, Lakshmanan R: The FOUR score predicts outcome in patients after traumatic brain injury. *Neurocrit Care* 2012; 16:95–101
66. Maas AI, Steyerberg EW, Butcher I, et al: Prognostic value of computerized tomography scan characteristics in traumatic brain injury: Results from the IMPACT study. *J Neurotrauma* 2007; 24:303–314
67. Maas AI, Hukkelhoven CW, Marshall LF, et al: Prediction of outcome in traumatic brain injury with computed tomographic characteristics: A comparison between the computed tomographic classification and combinations of computed tomographic predictors. *Neurosurgery* 2005; 57:1173–1182
68. Marshall LF, Toole BM, Bowers SA: The National Traumatic Coma Data Bank. Part 2: Patients who talk and deteriorate: Implications for treatment. *J Neurosurg* 1983; 59:285–288
69. Marino R, Gasparotti R, Pinelli L, et al: Posttraumatic cerebral infarction in patients with moderate or severe head trauma. *Neurology* 2006; 67:1165–1171
70. Mittl RL, Grossman RI, Hiehle JF, et al: Prevalence of MR evidence of diffuse axonal injury in patients with mild head injury and normal head CT findings. *AJNR Am J Neuroradiol* 1994; 15:1583–1589
71. Duckworth JL, Stevens RD: Imaging brain trauma. *Curr Opin Crit Care* 2010; 16:92–97
72. Huisman TA, Schwamm LH, Schaefer PW, et al: Diffusion tensor imaging as potential biomarker of white matter injury in diffuse axonal injury. *AJNR Am J Neuroradiol* 2004; 25:370–376
73. Tong KA, Ashwal S, Holshouser BA, et al: Diffuse axonal injury in children: Clinical correlation with hemorrhagic lesions. *Ann Neurol* 2004; 56:36–50
74. Schaefer PW, Huisman TA, Sorensen AG, et al: Diffusion-weighted MR imaging in closed head injury: High correlation with initial glasgow coma scale score and score on modified Rankin scale at discharge. *Radiology* 2004; 233:58–66
75. Newcombe VF, Williams GB, Nortje J, et al: Analysis of acute traumatic axonal injury using diffusion tensor imaging. *Br J Neurosurg* 2007; 21:340–348
76. Tollard E, Galanaud D, Perlberg V, et al: Experience of diffusion tensor imaging and 1H spectroscopy for outcome prediction in severe traumatic brain injury: Preliminary results. *Crit Care Med* 2009; 37:1448–1455
77. Sidaros A, Engberg AW, Sidaros K, et al: Diffusion tensor imaging during recovery from severe traumatic brain injury and relation to clinical outcome: A longitudinal study. *Brain* 2008; 131(Pt 2):559–572
78. Galanaud D, Perlberg V, Gupta R, et al: Assessment of white matter injury and outcome in severe brain trauma: A prospective multicenter cohort. *Anesthesiology* 2012; 117:1300–1310
79. Garnett MR, Blamire AM, Corkill RG, et al: Early proton magnetic resonance spectroscopy in normal-appearing brain correlates with outcome in patients following traumatic brain injury. *Brain* 2000; 123(Pt 10):2046–2054
80. Marino S, Zei E, Battaglini M, et al: Acute metabolic brain changes following traumatic brain injury and their relevance to clinical severity and outcome. *J Neurol Neurosurg Psychiatry* 2007; 78:501–507
81. Vik A, Nag T, Fredrikli OA, et al: Relationship of “dose” of intracranial hypertension to outcome in severe traumatic brain injury. *J Neurosurg* 2008; 109:678–684
82. Hiler M, Czosnyka M, Hutchinson P, et al: Predictive value of initial computerized tomography scan, intracranial pressure, and state of autoregulation in patients with traumatic brain injury. *J Neurosurg* 2006; 104:731–737
83. O’Sullivan MG, Statham PF, Jones PA, et al: Role of intracranial pressure monitoring in severely head-injured patients without signs of intracranial hypertension on initial computerized tomography. *J Neurosurg* 1994; 80:46–50
84. Juul N, Morris GF, Marshall SB, et al: Intracranial hypertension and cerebral perfusion pressure: Influence on neurological deterioration and outcome in severe head injury. The Executive Committee of the International Selfotel Trial. *J Neurosurg* 2000; 92:1–6
85. Brain Trauma Foundation: Guidelines for the management of severe traumatic brain injury. *J Neurotrauma* 2007; 24(Suppl 1):S1–106
86. Cremer OL, van Dijk GW, van Wensen E, et al: Effect of intracranial pressure monitoring and targeted intensive care on functional outcome after severe head injury. *Crit Care Med* 2005; 33:2207–2213
87. Shafi S, Diaz-Arrastia R, Madden C, et al: Intracranial pressure monitoring in brain-injured patients is associated with worsening of survival. *J Trauma* 2008; 64:335–340
88. Chesnut RM, Temkin N, Carney N, et al: A trial of intracranial-pressure monitoring in traumatic brain injury. *N Engl J Med* 2012; 367:2471–2481
89. Stiefel MF, Udoetuk JD, Spiotta AM, et al: Conventional neurocritical care and cerebral oxygenation after traumatic brain injury. *J Neurosurg* 2006; 105:568–575
90. Chang JJ, Youn TS, Benson D, et al: Physiologic and functional outcome correlates of brain tissue hypoxia in traumatic brain injury. *Crit Care Med* 2009; 37:283–290
91. Spiotta AM, Stiefel MF, Gracias VH, et al: Brain tissue oxygen-directed management and outcome in patients with severe traumatic brain injury. *J Neurosurg* 2010; 113:571–580
92. Stiefel MF, Spiotta A, Gracias VH, et al: Reduced mortality rate in patients with severe traumatic brain injury treated with brain tissue oxygen monitoring. *J Neurosurg* 2005; 103:805–811
93. Narotam PK, Morrison JF, Nathoo N: Brain tissue oxygen monitoring in traumatic brain injury and major trauma: Outcome analysis of a brain tissue oxygen-directed therapy. *J Neurosurg* 2009; 111:672–682
94. Vespa P, Bergsneider M, Hattori N, et al: Metabolic crisis without brain ischemia is common after traumatic brain injury: A combined microdialysis and positron emission tomography study. *J Cereb Blood Flow Metab* 2005; 25:763–774
95. Engström M, Polito A, Reinstrup P, et al: Intracerebral microdialysis in severe brain trauma: The importance of catheter location. *J Neurosurg* 2005; 102:460–469
96. Timofeev I, Carpenter KL, Nortje J, et al: Cerebral extracellular chemistry and outcome following traumatic brain injury: A microdialysis study of 223 patients. *Brain* 2011; 134(Pt 2):484–494
97. Marcoux J, McArthur DA, Miller C, et al: Persistent metabolic crisis as measured by elevated cerebral microdialysis lactate-pyruvate ratio predicts chronic frontal lobe brain atrophy after traumatic brain injury. *Crit Care Med* 2008; 36:2871–2877
98. Petzold A, Tisdall MM, Girbes AR, et al: In vivo monitoring of neuronal loss in traumatic brain injury: A microdialysis study. *Brain* 2011; 134(Pt 2):464–483
99. Magnoni S, Esparza TJ, Conte V, et al: Tau elevations in the brain extracellular space correlate with reduced amyloid-beta levels and predict adverse clinical outcomes after severe traumatic brain injury. *Brain* 2011; 135(Pt 4):1268–1280
100. Brody DL, Magnoni S, Schwetkey KE, et al: Amyloid-beta dynamics correlate with neurological status in the injured human brain. *Science* 2008; 321:1221–1224
101. Ronne-Engstrom E, Winkler T: Continuous EEG monitoring in patients with traumatic brain injury reveals a high incidence of epileptiform activity. *Acta Neurol Scand* 2006; 114:47–53
102. Wiedemayer H, Triesch K, Schäfer H, et al: Early seizures following non-penetrating traumatic brain injury in adults: Risk factors and clinical significance. *Brain Inj* 2002; 16:323–330
103. Vespa PM, Nuwer MR, Nenov V, et al: Increased incidence and impact of nonconvulsive and convulsive seizures after traumatic brain injury as detected by continuous electroencephalographic monitoring. *J Neurosurg* 1999; 91:750–760

104. Vespa PM, Miller C, McArthur D, et al: Nonconvulsive electrographic seizures after traumatic brain injury result in a delayed, prolonged increase in intracranial pressure and metabolic crisis. *Crit Care Med* 2007; 35:2830–2836
105. Vespa PM, McArthur DL, Xu Y, et al: Nonconvulsive seizures after traumatic brain injury are associated with hippocampal atrophy. *Neurology* 2010; 75:792–798
106. Brain Trauma Foundation. Guidelines for the management of severe traumatic brain injury. XIII. Antiseizure prophylaxis. *J Neurotrauma* 2007; 24(Suppl 1):S83–S86
107. Carter BG, Butt W: Review of the use of somatosensory evoked potentials in the prediction of outcome after severe brain injury. *Crit Care Med* 2001; 29:178–186
108. Houlden DA, Taylor AB, Feinstein A, et al: Early somatosensory evoked potential grades in comatose traumatic brain injury patients predict cognitive and functional outcome. *Crit Care Med* 2010; 38:167–174
109. Naeimi ZS, Weinhofer A, Sarahrudi K, et al: Predictive value of S-100B protein and neuron specific-enolase as markers of traumatic brain damage in clinical use. *Brain Inj* 2006; 20:463–468
110. Romner B, Ingebrigtsen T, Kongstad P, et al: Traumatic brain damage: Serum S-100 protein measurements related to neuroradiological findings. *J Neurotrauma* 2000; 17:641–647
111. Olivecrona M, Rodling-Wahlström M, Naredi S, et al: S-100B and neuron specific enolase are poor outcome predictors in severe traumatic brain injury treated by an intracranial pressure targeted therapy. *J Neurol Neurosurg Psychiatry* 2009; 80:1241–1247
112. Vos PE, Jacobs B, Andriessen TM, et al: GFAP and S100B are biomarkers of traumatic brain injury: An observational cohort study. *Neurology* 2010; 75:1786–1793
113. Rainey T, Lesko M, Sacho R, et al: Predicting outcome after severe traumatic brain injury using the serum S100B biomarker: Results using a single (24h) time-point. *Resuscitation* 2009; 80:341–345
114. Korfiatis S, Stranjalis G, Boviatisis E, et al: Serum S-100B protein monitoring in patients with severe traumatic brain injury. *Intensive Care Med* 2007; 33:255–260
115. Nylén K, Ost M, Csajbok LZ, et al: Serum levels of S100B, S100A1B and S100BB are all related to outcome after severe traumatic brain injury. *Acta Neurochir (Wien)* 2008; 150:221–227
116. Wiesmann M, Steinmeier E, Magerkurth O, et al: Outcome prediction in traumatic brain injury: Comparison of neurological status, CT findings, and blood levels of S100B and GFAP. *Acta Neurol Scand* 2010; 121:178–185
117. Böhmer AE, Oses JP, Schmidt AP, et al: Neuron-specific enolase, S100B, and glial fibrillary acidic protein levels as outcome predictors in patients with severe traumatic brain injury. *Neurosurgery* 2011; 68:1624–1630
118. Topolovec-Vranic J, Pollmann-Mudryj MA, Ouchterlony D, et al: The value of serum biomarkers in prediction models of outcome after mild traumatic brain injury. *J Trauma* 2011; 71(5 Suppl 1):S478–S486
119. Vos PE, Lamers KJ, Hendriks JC, et al: Glial and neuronal proteins in serum predict outcome after severe traumatic brain injury. *Neurology* 2004; 62:1303–1310
120. Pelinka LE, Kroepfl A, Leixnering M, et al: GFAP versus S100B in serum after traumatic brain injury: Relationship to brain damage and outcome. *J Neurotrauma* 2004; 21:1553–1561
121. Nylén K, Ost M, Csajbok LZ, et al: Increased serum-GFAP in patients with severe traumatic brain injury is related to outcome. *J Neurol Sci* 2006; 240:85–91
122. Liliang PC, Liang CL, Lu K, et al: Relationship between injury severity and serum tau protein levels in traumatic brain injured rats. *Resuscitation* 2010; 81:1205–1208
123. Liliang PC, Liang CL, Weng HC, et al: Tau proteins in serum predict outcome after severe traumatic brain injury. *J Surg Res* 2010; 160:302–307
124. Van Beek JG, Mushkudiani NA, Steyerberg EW, et al: Prognostic value of admission laboratory parameters in traumatic brain injury: Results from the IMPACT study. *J Neurotrauma* 2007; 24:315–328
125. Saggari V, Mittal RS, Vyas MC: Hemostatic abnormalities in patients with closed head injuries and their role in predicting early mortality. *J Neurotrauma* 2009; 26:1665–1668
126. Oddo M, Levine JM, Kumar M, et al: Anemia and brain oxygen after severe traumatic brain injury. *Intensive Care Med* 2012; 38:1497–1504
127. Nelson DW, Rudehill A, MacCallum RM, et al: Multivariate outcome prediction in traumatic brain injury with focus on laboratory values. *J Neurotrauma* 2012; 29:2613–2624
128. Greuters S, van den Berg A, Franschman G, et al: Acute and delayed mild coagulopathy are related to outcome in patients with isolated traumatic brain injury. *Crit Care* 2011; 15:R2
129. Jeremitsky E, Omert LA, Dunham CM, et al: The impact of hyperglycemia on patients with severe brain injury. *J Trauma* 2005; 58:47–50
130. Bilotta F, Caramia R, Cernak I, et al: Intensive insulin therapy after severe traumatic brain injury: A randomized clinical trial. *Neurocrit Care* 2008; 9:159–166
131. Yang M, Guo Q, Zhang X, et al: Intensive insulin therapy on infection rate, days in NICU, in-hospital mortality and neurological outcome in severe traumatic brain injury patients: A randomized controlled trial. *Int J Nurs Stud* 2009; 46:753–758
132. Green DM, O'Phelan KH, Bassin SL, et al: Intensive versus conventional insulin therapy in critically ill neurologic patients. *Neurocrit Care* 2010; 13:299–306
133. Vespa P, Boonyaputthikul R, McArthur DL, et al: Intensive insulin therapy reduces microdialysis glucose values without altering glucose utilization or improving the lactate/pyruvate ratio after traumatic brain injury. *Crit Care Med* 2006; 34:850–856
134. Meierhans R, Béchir M, Ludwig S, et al: Brain metabolism is significantly impaired at blood glucose below 6mM and brain glucose below 1mM in patients with severe traumatic brain injury. *Crit Care* 2010; 14:R13
135. Béchir M, Meierhans R, Brandi G, et al: Insulin differentially influences brain glucose and lactate in traumatic brain injured patients. *Minerva Anesthesiol* 2010; 76:896–904
136. Vespa P, McArthur DL, Stein N, et al: Tight glycemic control increases metabolic distress in traumatic brain injury: A randomized controlled within-subjects trial. *Crit Care Med* 2012; 40:1923–1929
137. Andrews PJ, Sleeman DH, Statham PF, et al: Predicting recovery in patients suffering from traumatic brain injury by using admission variables and physiological data: A comparison between decision tree analysis and logistic regression. *J Neurosurg* 2002; 97:326–336
138. Clifton GL, Miller ER, Choi SC, et al: Fluid thresholds and outcome from severe brain injury. *Crit Care Med* 2002; 30:739–745
139. Davis DP, Peay J, Sise MJ, et al: Prehospital airway and ventilation management: A trauma score and injury severity score-based analysis. *J Trauma* 2010; 69:294–301
140. Bulger EM, Nathens AB, Rivara FP, et al: Management of severe head injury: Institutional variations in care and effect on outcome. *Crit Care Med* 2002; 30:1870–1876
141. Patel HC, Menon DK, Tebbs S, et al: Specialist neurocritical care and outcome from head injury. *Intensive Care Med* 2002; 28:547–553
142. Clifton GL, Valadka A, Zygun D, et al: Very early hypothermia induction in patients with severe brain injury (the National Acute Brain Injury Study: Hypothermia II): A randomised trial. *Lancet Neurol* 2011; 10:131–139
143. Clifton GL, Miller ER, Choi SC, et al: Lack of effect of induction of hypothermia after acute brain injury. *N Engl J Med* 2001; 344:556–563
144. Sydenham E, Roberts I, Alderson P. Hypothermia for traumatic head injury. *Cochr Database Syst Rev* 2009; (1):CD001048
145. Wijdicks EF, Hijdra A, Young GB, et al: Practice parameter: Prediction of outcome in comatose survivors after cardiopulmonary resuscitation (an evidence-based review): Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2006; 67:203–210
146. Zandbergen EG, Hijdra A, Koelman JH, et al: Prediction of poor outcome within the first 3 days of postanoxic coma. *Neurology* 2006; 66:62–68
147. Bates D, Caronna JJ, Carlidge NE, et al: A prospective study of

- nontraumatic coma: Methods and results in 310 patients. *Ann Neurol* 1977; 2:211–220
148. Levy DE, Caronna JJ, Singer BH, et al: Predicting outcome from hypoxic-ischemic coma. *JAMA* 1985; 253:1420–1426
 149. Greer DM, Yang J, Scripko PD, et al: Clinical examination for outcome prediction in nontraumatic coma. *Crit Care Med* 2012; 40:1150–1156
 150. Bouwes A, Binnekade JM, Verbaan BW, et al: Predictive value of neurological examination for early cortical responses to somatosensory evoked potentials in patients with postanoxic coma. *J Neurol* 2012; 259:537–541
 151. Rossetti AO, Oddo M, Logroscino G, et al: Prognostication after cardiac arrest and hypothermia: A prospective study. *Ann Neurol* 2010; 67:301–307
 152. Posner JB, Plum F, Saper CB. Plum and Posner's Diagnosis of Stupor and Coma. Oxford, Oxford University Press, 2007
 153. Inamasu J, Miyatake S, Nakatsukasa M, et al: Loss of gray-white matter discrimination as an early CT sign of brain ischemia/hypoxia in victims of asphyxial cardiac arrest. *Emerg Radiol* 2011; 18:295–298
 154. Torbey MT, Selim M, Knorr J, et al: Quantitative analysis of the loss of distinction between gray and white matter in comatose patients after cardiac arrest. *Stroke* 2000; 31:2163–2167
 155. Wijndicks EF, Campeau NG, Miller GM: MR imaging in comatose survivors of cardiac resuscitation. *AJNR Am J Neuroradiol* 2001; 22:1561–1565
 156. Chalela JA, Wolf RL, Maldjian JA, et al: MRI identification of early white matter injury in anoxic-ischemic encephalopathy. *Neurology* 2001; 56:481–485
 157. Singhal AB, Topcuoglu MA, Koroshetz WJ: Diffusion MRI in three types of anoxic encephalopathy. *J Neurol Sci* 2002; 196:37–40
 158. Wu O, Sorensen AG, Benner T, et al: Comatose patients with cardiac arrest: Predicting clinical outcome with diffusion-weighted MR imaging. *Radiology* 2009; 252:173–181
 159. Wijman CA, Mlynash M, Caulfield AF, et al: Prognostic value of brain diffusion-weighted imaging after cardiac arrest. *Ann Neurol* 2009; 65:394–402
 160. Choi SP, Park KN, Park HK, et al: Diffusion-weighted magnetic resonance imaging for predicting the clinical outcome of comatose survivors after cardiac arrest: A cohort study. *Crit Care* 2010; 14:R17
 161. Mlynash M, Campbell DM, Leproust EM, et al: Temporal and spatial profile of brain diffusion-weighted MRI after cardiac arrest. *Stroke* 2010; 41:1665–1672
 162. Luyt CE, Galanaud D, Perlberg V, et al: Diffusion tensor imaging to predict long-term outcome following cardiac arrest: A bi-centric pilot study. *Anesthesiology* 2012; 117(6):1311–1321
 163. Graham SH, Meyerhoff DJ, Bayne L, et al: Magnetic resonance spectroscopy of N-acetylaspartate in hypoxic-ischemic encephalopathy. *Ann Neurol* 1994; 35:490–494
 164. Wartenberg KE, Patsalides A, Yepes MS: Is magnetic resonance spectroscopy superior to conventional diagnostic tools in hypoxic-ischemic encephalopathy? *J Neuroimaging* 2004; 14:180–186
 165. Krumholz A, Stern BJ, Weiss HD: Outcome from coma after cardiopulmonary resuscitation: Relation to seizures and myoclonus. *Neurology* 1988; 38:401–405
 166. Snyder BD, Hauser WA, Loewenson RB, et al: Neurologic prognosis after cardiopulmonary arrest: III. Seizure activity. *Neurology* 1980; 30:1292–1297
 167. Chen R, Bolton CF, Young B: Prediction of outcome in patients with anoxic coma: A clinical and electrophysiologic study. *Crit Care Med* 1996; 24:672–678
 168. Rossetti AO, Oddo M, Liaudet L, et al: Predictors of awakening from postanoxic status epilepticus after therapeutic hypothermia. *Neurology* 2009; 72:744–749
 169. Geocadin RG, Buitrago MM, Torbey MT, et al: Neurologic prognosis and withdrawal of life support after resuscitation from cardiac arrest. *Neurology* 2006; 67:105–108
 170. Wennervirta JE, Ermes MJ, Tiainen SM, et al: Hypothermia-treated cardiac arrest patients with good neurological outcome differ early in quantitative variables of EEG suppression and epileptiform activity. *Crit Care Med* 2009; 37:2427–2435
 171. Rossetti AO, Urbano LA, Delodder F, et al: Prognostic value of continuous EEG monitoring during therapeutic hypothermia after cardiac arrest. *Crit Care* 2010; 14:R173
 172. Robinson LR, Micklesen PJ, Tirschwell DL, et al: Predictive value of somatosensory evoked potentials for awakening from coma. *Crit Care Med* 2003; 31:960–967
 173. Zandbergen EG, Koelman JH, de Haan RJ, et al: SSEPs and prognosis in postanoxic coma: Only short or also long latency responses? *Neurology* 2006; 67:583–586
 174. Lee YC, Phan TG, Jolley DJ, et al: Accuracy of clinical signs, SEP, and EEG in predicting outcome of hypoxic coma: A meta-analysis. *Neurology* 2010; 74:572–580
 175. Kane NM, Butler SR, Simpson T: Coma outcome prediction using event-related potentials: P(3) and mismatch negativity. *Audiol Neurootol* 2000; 5:186–191
 176. Young GB, Wang JT, Connolly JF: Prognostic determination in anoxic-ischemic and traumatic encephalopathies. *J Clin Neurophysiol* 2004; 21:379–390
 177. Mecklinger A, Opitz B, Friederici AD: Semantic aspects of novelty detection in humans. *Neurosci Lett* 1997; 235:65–68
 178. Fischer C, Luauté J, Némaz C, et al: Improved prediction of awakening or nonawakening from severe anoxic coma using tree-based classification analysis. *Crit Care Med* 2006; 34:1520–1524
 179. Tiainen M, Poutiainen E, Kovala T, et al: Cognitive and neurophysiological outcome of cardiac arrest survivors treated with therapeutic hypothermia. *Stroke* 2007; 38:2303–2308
 180. Vaagenes P, Kjekshus J, Torvik A: The relationship between cerebrospinal fluid creatine kinase and morphologic changes in the brain after transient cardiac arrest. *Circulation* 1980; 61:1194–1199
 181. Kärkelä J, Pasanen M, Kaukinen S, et al: Evaluation of hypoxic brain injury with spinal fluid enzymes, lactate, and pyruvate. *Crit Care Med* 1992; 20:378–386
 182. Müllner M, Sterz F, Domanovits H, et al: The association between blood lactate concentration on admission, duration of cardiac arrest, and functional neurological recovery in patients resuscitated from ventricular fibrillation. *Intensive Care Med* 1997; 23:1138–1143
 183. Fogel W, Krieger D, Veith M, et al: Serum neuron-specific enolase as early predictor of outcome after cardiac arrest. *Crit Care Med* 1997; 25:1133–1138
 184. Rosén H, Rosengren L, Herlitz J, et al: Increased serum levels of the S-100 protein are associated with hypoxic brain damage after cardiac arrest. *Stroke* 1998; 29:473–477
 185. Meynaar IA, Oudemans-van Straaten HM, van der Wetering J, et al: Serum neuron-specific enolase predicts outcome in post-anoxic coma: A prospective cohort study. *Intensive Care Med* 2003; 29:189–195
 186. Tiainen M, Roine RO, Pettilä V, et al: Serum neuron-specific enolase and S-100B protein in cardiac arrest patients treated with hypothermia. *Stroke* 2003; 34:2881–2886
 187. Fugate JE, Wijndicks EF, Mandrekar J, et al: Predictors of neurologic outcome in hypothermia after cardiac arrest. *Ann Neurol* 2010; 68:907–914
 188. Steffen IG, Hasper D, Ploner CJ, et al: Mild therapeutic hypothermia alters neuron specific enolase as an outcome predictor after resuscitation: 97 prospective hypothermia patients compared to 133 historical non-hypothermia patients. *Crit Care* 2010; 14:R69
 189. Bouwes A, Binnekade JM, Kuiper MA, et al: Prognosis of coma after therapeutic hypothermia: A prospective cohort study. *Ann Neurol* 2012; 71:206–212
 190. Rossetti AO, Carrera E, Oddo M: Early EEG correlates of neuronal injury after brain anoxia. *Neurology* 2012; 78:796–802
 191. Kaneko T, Kasaoka S, Miyauchi T, et al: Serum glial fibrillary acidic protein as a predictive biomarker of neurological outcome after cardiac arrest. *Resuscitation* 2009; 80:790–794
 192. Hayashida H, Kaneko T, Kasaoka S, et al: Comparison of the predictability of neurological outcome by serum procalcitonin and glial fibrillary acidic protein in postcardiac-arrest patients. *Neurocrit Care* 2010; 12:252–257
 193. Randall J, Lord B, Fulham J, et al: Parastomal hernias as the predominant stoma complication after laparoscopic

- colorectal surgery. *Surg Laparosc Endosc Percutan Tech* 2012; 22: 420–423
194. Zeiner A, Holzer M, Sterz F, et al: Hyperthermia after cardiac arrest is associated with an unfavorable neurologic outcome. *Arch Intern Med* 2001; 161:2007–2012
 195. Wolff B, Machill K, Schumacher D, et al: Early achievement of mild therapeutic hypothermia and the neurologic outcome after cardiac arrest. *Int J Cardiol* 2009; 133:223–228
 196. Bernard SA, Smith K, Cameron P, et al: Induction of therapeutic hypothermia by paramedics after resuscitation from out-of-hospital ventricular fibrillation cardiac arrest: A randomized controlled trial. *Circulation* 2010; 122:737–742
 197. Castrén M, Nordberg P, Svensson L, et al: Intra-arrest transnasal evaporative cooling: A randomized, prehospital, multicenter study (PRINCE: Pre-ROSC IntraNasal Cooling Effectiveness). *Circulation* 2010; 122:729–736
 198. De Georgia M, Raad B: Prognosis of coma after cardiac arrest in the era of hypothermia. *Continuum (Minneap Minn)* 2012; 18: 515–531
 199. Zandbergen EG, de Haan RJ, Stoutenbeek CP, et al: Systematic review of early prediction of poor outcome in anoxic-ischaemic coma. *Lancet* 1998; 352:1808–1812
 200. Booth CM, Boone RH, Tomlinson G, et al: Is this patient dead, vegetative, or severely neurologically impaired? Assessing outcome for comatose survivors of cardiac arrest. *JAMA* 2004; 291: 870–879
 201. Al Thenayan E, Savard M, Sharpe M, et al: Predictors of poor neurologic outcome after induced mild hypothermia following cardiac arrest. *Neurology* 2008; 71:1535–1537
 202. Rossetti AO, Oddo M, Logroscino G, et al: Prognostication after cardiac arrest and hypothermia: A prospective study. *Ann Neurol* 2010; 5(2):161–174
 203. Wijdicks EF, Parisi JE, Sharbrough FW: Prognostic value of myoclonus status in comatose survivors of cardiac arrest. *Ann Neurol* 1994; 35:239–243
 204. Zandbergen EG, de Haan RJ, Hijdra A: Systematic review of prediction of poor outcome in anoxic-ischaemic coma with biochemical markers of brain damage. *Intensive Care Med* 2001; 27:1661–1667
 205. Zingler VC, Krumm B, Bertsch T, et al: Early prediction of neurological outcome after cardiopulmonary resuscitation: A multimodal approach combining neurobiochemical and electrophysiological investigations may provide high prognostic certainty in patients after cardiac arrest. *Eur Neurol* 2003; 49:79–84
 206. Cronberg T, Rundgren M, Westhall E, et al: Neuron-specific enolase correlates with other prognostic markers after cardiac arrest. *Neurology* 2011; 77:623–630
 207. Pohlmann-Eden B, Dingethal K, Bender HJ, et al: How reliable is the predictive value of SEP (somatosensory evoked potentials) patterns in severe brain damage with special regard to the bilateral loss of cortical responses? *Intensive Care Med* 1997; 23:301–308
 208. Bouwes A, Binnekade JM, Zandstra DF, et al: Somatosensory evoked potentials during mild hypothermia after cardiopulmonary resuscitation. *Neurology* 2009; 73:1457–1461
 209. Leithner C, Ploner CJ, Hasper D, et al: Does hypothermia influence the predictive value of bilateral absent N20 after cardiac arrest? *Neurology* 2010; 74:965–969
 210. Rossetti AO, Logroscino G, Liaudet L, et al: Status epilepticus: An independent outcome predictor after cerebral anoxia. *Neurology* 2007; 69:255–260
 211. Thenayan EA, Savard M, Sharpe MD, et al: Electroencephalogram for prognosis after cardiac arrest. *J Crit Care* 2010; 25:300–304
 212. Howard RS, Holmes PA, Siddiqui A, et al: Hypoxic-ischaemic brain injury: Imaging and neurophysiology abnormalities related to outcome. *QJM* 2012; 105:551–561
 213. Topcuoglu MA, Oguz KK, Buyukserbetci G, et al: Prognostic value of magnetic resonance imaging in post-resuscitation encephalopathy. *Intern Med* 2009; 48:1635–1645