

## REVIEW ARTICLE

Jeffrey M. Drazen, M.D., *Editor*

## Acute Respiratory Distress Syndrome

B. Taylor Thompson, M.D., Rachel C. Chambers, Ph.D., and Kathleen D. Liu, M.D., Ph.D.

From the Division of Pulmonary and Critical Care, Department of Medicine, Massachusetts General Hospital, and Harvard Medical School — both in Boston (B.T.T.); Centre for Inflammation and Tissue Repair, the Division of Medicine, University College London, London (R.C.C.); and the Divisions of Nephrology and Critical Care Medicine, University of California San Francisco, San Francisco (K.D.L.). Address reprint requests to Dr. Thompson at the Division of Pulmonary and Critical Care, Department of Medicine, Massachusetts General Hospital, Bulfinch Bldg., Suite 148, 55 Fruit St., Boston, MA 02114, or at thompson.taylor@mgh.harvard.edu.

This article was last updated on August 11, 2017, at NEJM.org.

N Engl J Med 2017;377:562-72.

DOI: 10.1056/NEJMra1608077

Copyright © 2017 Massachusetts Medical Society.

FIFTY YEARS AGO, ASHBAUGH AND COLLEAGUES DESCRIBED 12 PATIENTS WITH tachypnea, refractory hypoxemia, and diffuse opacities on chest radiographs after infection or trauma.<sup>1</sup> Prominent hyaline membranes were seen lining the alveolar spaces of the lungs in 6 of the 7 patients who died, findings previously thought to be specific for the respiratory distress syndrome of the newborn. Thus, the term adult (later changed to acute) respiratory distress syndrome (ARDS) was proposed.

Since ARDS was last reviewed in the *Journal*, 17 years ago,<sup>2</sup> substantial progress has been made in the care of affected patients and those at risk for the disorder, with reductions in both incidence and mortality. However, ARDS remains a relatively common and lethal or disabling syndrome. In a recent international study involving 29,144 patients,<sup>3</sup> 10% of all patients admitted to the intensive care unit (ICU) and 23% of mechanically ventilated patients had ARDS. Mortality in the subgroup of patients with severe ARDS was 46%.<sup>3</sup> Patients who survive this disorder are at high risk for cognitive decline, depression, post-traumatic stress disorder, and persistent skeletal-muscle weakness.<sup>4,5</sup>

## DEFINITION AND PATHOLOGICAL FEATURES

Four major definitions of ARDS have evolved over the years, and all have retained the central features of the initial description by Ashbaugh and colleagues. Because lung permeability, edema, and inflammation are not routinely measured in clinical care and no validated diagnostic biomarkers are yet available, these definitions rely on clinical features and chest imaging as surrogates. The Berlin definition, proposed in 2012,<sup>6</sup> breaks with tradition by establishing three risk strata that are based on the degree of hypoxemia as assessed at a minimum positive end-expiratory pressure (PEEP) (Table 1). The definition makes the radiographic criteria more explicit and allows the use of computed tomography (CT) for the detection of qualifying opacities, which are often heterogeneous (Fig. 1). In addition, the definition acknowledges that if ARDS develops, it usually does so within 7 days after clinical recognition of a known risk factor, most commonly pneumonia or sepsis (Table 2). ARDS with a more indolent onset or in the absence of an identifiable risk factor should prompt consideration of so-called ARDS mimics, a large number of diseases or syndromes that may require specific treatments (Table 3).<sup>8</sup> Prior definitions excluded volume overload or heart failure, but recent evidence suggests that these problems may coexist in up to a third of patients with ARDS.

The histologic correlate of ARDS is widely considered to be “diffuse alveolar damage,” a term coined by Katzenstein and colleagues<sup>9</sup> almost a decade after the report by Ashbaugh et al.<sup>1</sup> Katzenstein and colleagues described the rapid development of capillary congestion, atelectasis, intraalveolar hemorrhage, and alveolar edema, followed days later by hyaline-membrane formation, epithelial-cell hyperplasia, and interstitial edema. Animal models of ARDS have been developed in an effort to recapitulate these histologic findings. However, the Berlin definition (as well as the 1994 American–European Consensus Conference definition<sup>10</sup>) has poor specificity for diffuse alveolar damage. At postmortem examination, 40 to 58% of patients

**Table 1. Berlin Definition of the Acute Respiratory Distress Syndrome (ARDS).\***

Criteria	Rationale
Onset within 7 days after a known clinical insult or new or worsening respiratory symptoms	Observational data suggest that ARDS will develop within 72 hr in the majority of patients at risk for the syndrome and within 1 wk in nearly all patients at risk
Bilateral opacities that are “consistent with pulmonary edema” on chest radiographs or chest CT	There is poor interobserver reliability in interpreting the chest radiograph for the presence of edema. To address this issue, the Berlin definition offers more explicit criteria (e.g., opacities should not be fully explained by effusions, lobar or lung atelectasis, or nodules or masses), with illustrative radiographs provided
Categorization of ARDS severity	A patient-level meta-analysis validated three thresholds for hypoxemia, all consisting of a $P_{aO_2}:F_{iO_2}$ ratio $\leq 300$ mm Hg
Mild	$P_{aO_2}:F_{iO_2}$ , 201 to 300 mm Hg; mortality, 27% (95% CI, 24–30)
Moderate	$P_{aO_2}:F_{iO_2}$ , 101 to 200 mm Hg; mortality, 32% (95% CI, 29–34)
Severe	$P_{aO_2}:F_{iO_2}$ , $\leq 100$ mm Hg; mortality, 45% (95% CI, 42–48)
Minimum PEEP setting or CPAP, 5 cm of water; $P_{aO_2}:F_{iO_2}$ assessed on invasive mechanical ventilation (CPAP criterion used for the diagnosis of mild ARDS)	Estimates of $F_{iO_2}$ are not accurate with oxygen-delivery systems other than invasive or non-invasive ventilation (with a tight-fitting mask), with the exception of nasal high-flow oxygen delivery systems (at flow rates $\geq 45$ liters per minute); requiring higher PEEP settings does not increase predictive validity of the Berlin severity strata and adds complexity

\* The definition and the quotation about opacities are from Ferguson et al.<sup>6</sup> CI denotes confidence interval, CPAP continuous positive airway pressure,  $P_{aO_2}:F_{iO_2}$  ratio of the partial pressure of arterial oxygen to the fraction of inspired oxygen, and PEEP positive end-expiratory pressure.

with a clinical diagnosis of moderate-to-severe ARDS have diffuse alveolar damage. Pulmonary edema and pneumonia without hyaline membranes are the next most common findings, although 14% of patients have no pulmonary lesions whatsoever, probably because of atelectasis masquerading as ARDS.<sup>11,12</sup> Similar or lower proportions of patients have diffuse alveolar damage on lung biopsy. Furthermore, the overall proportion of patients with diffuse alveolar damage at postmortem examination has fallen from 49% to 41% in the past decade, as mechanical ventilation with tidal volumes on the order of 6 ml per kilogram of ideal body weight has become common.<sup>12</sup> Thus, diffuse alveolar damage is best thought of as a common histologic finding in patients with ARDS that may, in part, reflect ventilator-induced lung injury.

#### EPIDEMIOLOGIC FEATURES


Population-based estimates of ARDS range from 10 to 86 cases per 100,000, with the highest rates reported in Australia and the United States.<sup>13</sup> ARDS is likely to be underreported in low-income countries, where resources to obtain chest radiographs and measure arterial blood gases are limited. In Kigali, Rwanda, for example, Riviello and colleagues<sup>14</sup> found no cases of ARDS when they applied the Berlin definition, but with the substitution of lung ultrasonography for radiography or CT and percutaneous oxygen satura-

tion for blood gas measurements, they identified 4 cases per 1000 hospital admissions, with a 50% mortality rate.

Even in high-income countries, ARDS remains underrecognized. A recent observational study of 459 ICUs in 50 countries showed clinical recognition rates ranging from 51.3% for mild ARDS to 78.5% for severe ARDS.<sup>3</sup> Underuse of currently recommended lung-protective practices was seen across the severity spectrum, suggesting both underrecognition and undertreatment. One explanation for the underdiagnosis of ARDS may be disagreement about the nature of the radiographic opacities that support the diagnosis. Diffuse, confluent opacities with a narrow cardiothoracic silhouette (suggesting noncardiogenic edema) are the classic findings, but imaging often shows asymmetric, dependent, and occasionally lobar opacities (Fig. 1). Patients with multifocal opacities have been enrolled in ARDS clinical trials for decades and benefit from lung-protective ventilation.<sup>15</sup> Computerized detection algorithms reliably identify ARDS from the electronic medical record in real time and offer a solution to the problem of underrecognition at some tertiary care centers, but such algorithms have not been widely implemented or validated.<sup>16</sup>

#### GENETIC FEATURES AND BIOMARKERS

ARDS does not develop in the majority of patients with clinical risk factors for the disease



(e.g., pneumonia, sepsis, or trauma), suggesting that other factors, including genetic susceptibility, play a key role in the pathogenesis of this disorder. However, differences in virulence fac-

tors (e.g., H1N1 influenza), coexisting conditions (e.g., pneumococcal pneumonia after splenectomy), and environmental exposures (alcohol use or active smoking and injurious mechanical-

**Figure 1 (facing page). Radiographic Heterogeneity in Patients with the Acute Respiratory Distress Syndrome (ARDS).**

Panels A through D are radiographs included with the publication of the Berlin definition<sup>6</sup> that meet the diagnostic criteria for ARDS. The radiographs and an explicit definition of qualifying opacities were published with the goal of reducing poor interobserver reliability. Qualifying opacities must be bilateral and “consistent with pulmonary edema”; furthermore, the opacities cannot be “fully explained by effusions, lobar/lung collapse, or nodules/masses on chest radiograph.”<sup>6</sup> The Berlin definition also recognizes the use of CT for the detection of qualifying opacities so that a CT scan can be substituted for the chest radiograph.<sup>6</sup> Panel E is a CT image showing the heterogeneous nature of the opacities often seen in patients with ARDS, as exemplified in Panels C and D, and Panel F is an <sup>18</sup>F-fluorodeoxyglucose (FDG) positron-emission tomographic image corresponding to the CT image in Panel E.<sup>7</sup> Intense FDG uptake, largely from metabolically active inflammatory cells, is observed in normally aerated regions that receive a relatively larger fraction of the delivered tidal volume (rectangle 1 in Panels E and F), probably reflecting volutrauma. Activity is lower in the dorsal, nonaerated regions of both lungs (rectangle 2 in Panels E and F), in part reflecting atelectasis. All images have been reprinted from Ferguson et al.<sup>6</sup> and Bellani et al.<sup>7</sup> with the permission of the publishers.

**Table 2. Risk Factors for ARDS.****Direct lung-injury risk factors**

Pneumonia (bacterial, viral, fungal, or opportunistic)\*  
 Aspiration of gastric contents\*  
 Pulmonary contusion  
 Inhalation injury  
 Near drowning

**Indirect lung-injury risk factors**

Sepsis (nonpulmonary source)\*  
 Nonthoracic trauma or hemorrhagic shock  
 Pancreatitis  
 Major burn injury  
 Drug overdose  
 Transfusion of blood products  
 Cardiopulmonary bypass  
 Reperfusion edema after lung transplantation or embolectomy

\* Pneumonia, aspiration of gastric contents, and sepsis together account for more than 85% of cases of ARDS in recent clinical trials.

ventilation practices)<sup>17</sup> complicate the interpretation of genetic findings. More than 40 candidate genes associated with the development or outcome of ARDS have been identified, including the genes encoding angiotensin-converting enzyme (*ACE*), interleukin 10 (*IL-10*), tumor necrosis factor (*TNF*), and vascular endothelial growth factor (*VEGF*), as well as *SOD3*, *MYLK*, *NFE2L2*, *NAMPT*, and *SFTPB*.<sup>18</sup> In the one genomewide association study that has been reported for trauma-associated ARDS, no polymorphism had genomewide significance.<sup>19</sup> As with other diseases, the biologic importance of the genetic association is strengthened by additional studies that implicate the same pathway. For example, *ACE* has been associated with overall susceptibility to ARDS,<sup>20</sup> and the *ACE2* protein is the receptor for the severe acute respiratory syndrome coronavirus (SARS-CoV).<sup>21,22</sup> Experimentally induced lung injury from SARS-CoV can be attenuated by blocking the renin–angiotensin pathway,<sup>22,23</sup> suggesting both a molecular explanation for the severe ARDS that follows SARS-CoV infection and a possible treatment.

Sequencing of genomic coding regions (exome sequencing) identified polymorphisms in the genes encoding arylsulfatase D (*ARSD*) and X Kell

blood-group precursor–related family, member 3 (*XKR3*) and showed differences in expression between patients with ARDS and healthy controls, but these findings require replication.<sup>24</sup> Exome sequencing has also shown that more than one genetic variant may explain a clinical phenotype. Thus, some persons may have multiple variants that modify the risk of ARDS and the outcome of ARDS, which may go undetected or at least lead to imprecise risk estimates.<sup>25</sup> In addition, lung tissue for discovery research is generally not available from patients with ARDS, and even within the lung compartment, there are a number of different cell types that may not be cleanly separated. Thus, the overall success of linking candidate genes with ARDS susceptibility and outcomes, as well as with downstream biologic events (e.g., transcriptional and epigenetic events or protein expression), remains limited. Microengineered “lungs on a chip” and isolated perfused human lungs are two preclinical platforms that have the potential to bridge these gaps but have yet to be proved useful in identifying new treatments.<sup>26,27</sup>

Increased levels of plasma biomarkers, including markers of systemic inflammation (interleukin-6 and interleukin-8), epithelial injury (receptor for advanced glycation end products and surfactant protein D), and endothelial injury (angiopoietin 2), as well as markers of dysregu-

**Table 3. Conditions That May Mimic ARDS.\***

Congestive heart failure
Interstitial lung disease (e.g., acute interstitial pneumonia, nonspecific interstitial pneumonitis, cryptogenic organizing pneumonia, acute eosinophilic pneumonia, hypersensitivity pneumonia, and pulmonary alveolar proteinosis)
Connective-tissue diseases such as polymyositis (antisynthetase syndrome)
Diffuse alveolar hemorrhage from vasculitis or Goodpasture's syndrome
Drug-induced lung diseases (e.g., bleomycin or amiodarone), including vascular leak syndrome from immunotherapy
Cancer (T-cell or B-cell lymphomas or metastatic carcinoma)
Endobronchial tuberculosis

\* These conditions, referred to as "ARDS mimics" or "secondary causes" in the literature, may require additional diagnostic tests and treatments distinct from those for ARDS.<sup>8</sup>

lated coagulation (low protein C and high plasminogen activator inhibitor 1 levels), have been associated with adverse outcomes of ARDS. These biomarkers provide insights into the pathogenesis of ARDS and may identify treatment-responsive subtypes (see the Supplementary Appendix, available with the full text of this article at NEJM.org).

#### PATHOGENESIS

Figures 2 and 3 provide an overview of the pathogenesis of ARDS; a detailed review can be found in the Supplementary Appendix. The lung's initial response to injury, referred to as the exudative phase of ARDS, is characterized by innate immune cell-mediated damage of the alveolar endothelial and epithelial barriers and accumulation of protein-rich edema fluid within the interstitium and alveolus (Fig. 2). Resident alveolar macrophages secrete proinflammatory cytokines, leading to neutrophil and monocyte or macrophage recruitment, as well as activation of alveolar epithelial cells and effector T cells, to promote and sustain inflammation and tissue injury.<sup>29</sup> Endothelial activation and microvascular injury also contribute to the barrier disruption in ARDS and are worsened by mechanical stretch. The repair processes initiated during the second, or proliferative, phase of ARDS are essential for host survival (Fig. 3A). Once epithelial integrity has been reestablished, reabsorption of alveolar edema and the provisional matrix restores alveolar architecture and function. The final, or fibrotic, phase of ARDS (Fig. 3B) does not occur in all patients but has been linked to prolonged mechanical ventilation and increased mortality.

**Figure 2 (facing page). The Healthy Lung and the Exudative Phase of ARDS.**


The healthy lung is shown on the left, and the exudative phase of ARDS is shown on the right. Injury is initiated by either direct or indirect insults to the delicate alveolar structure of the distal lung and associated microvasculature. In the exudative phase, resident alveolar macrophages are activated, leading to the release of potent proinflammatory mediators and chemokines that promote the accumulation of neutrophils and monocytes. Activated neutrophils further contribute to injury by releasing toxic mediators. The resultant injury leads to loss of barrier function, as well as interstitial and intra-alveolar flooding. Tumor necrosis factor (TNF)-mediated expression of tissue factor promotes platelet aggregation and microthrombus formation, as well as intraalveolar coagulation and hyaline-membrane formation. AECI denotes type I alveolar epithelial cell, AECII type II alveolar epithelial cell, Ang-2 angiotensin-2, APC activated protein C, CC-16 club cell (formerly Clara cell) secretory protein 16, CCL chemokine (CC motif) ligand, DAMP damage-associated molecular pattern, ENaC epithelial sodium channel, GAG glycosaminoglycan, HMGB1 high-mobility group box 1 protein, KL-6 Krebs von den Lungen 6, LPS lipopolysaccharide, LTB4 leukotriene B4, MMP matrix metalloproteinase, MPO myeloperoxidase, mtDNA mitochondrial DNA, Na<sup>+</sup>/K<sup>+</sup> ATPase sodium-potassium ATPase pump, NF- $\kappa$ B nuclear factor kappa light-chain enhancer of activated B cells, NET neutrophil extracellular trap, PAMP pathogen-associated molecular pattern, PRR pattern recognition receptor, ROS reactive oxygen species, sICAM soluble intercellular adhesion molecule, SP surfactant protein, sRAGE soluble receptor for advanced glycation end products, VEGF vascular endothelial growth factor, and vWF von Willebrand factor.

#### TREATMENT AND PREVENTION

##### SUPPORTIVE THERAPY

The first priority in the care of patients with ARDS is identification and treatment of the underlying cause (or causes). For example, in patients with sepsis-associated ARDS, good outcomes require early resuscitation, appropriate antibiotic agents, and source control.<sup>30</sup>


Supportive therapy for ARDS is focused on limiting further lung injury through a combination of lung-protective ventilation to prevent ventilator-associated lung injury (reviewed in the *Journal* in 2013 by Slutsky and Ranieri<sup>31</sup>) and conservative fluid therapy to prevent lung edema formation and promote lung edema resorption. The optimal approach to lung-protective ventilation is unknown. Current evidence suggests that there may be no safe level of tidal volume or airway pressure in patients with acute lung in-



jury. Because the volume of aerated lung is reduced in patients with ARDS, even normal tidal volumes delivered with airway pressures that are considered safe for the uninjured lung may cause regional overdistention (so-called volutrauma), further activating or injuring the epithelium and amplifying inflammation. Repetitive opening and closing of lung units (atelectrauma) amplifies regional lung strain and denatures surfactant. Finally, epithelial and

endothelial injury results in translocation of proinflammatory mediators and bacterial products, leading to worsening systemic inflammation (biotrauma).

Clinical practice guidelines endorsed by multiple professional societies<sup>32</sup> recommend invasive mechanical ventilation after a lowering of the tidal volume and airway pressure. Tidal volumes are reduced from 6 ml per kilogram of predicted body weight to a minimum of 4 ml per kilogram



**Figure 3. The Proliferative and Fibrotic Phases of ARDS.**

The proliferative phase (Panel A) aims to restore tissue homeostasis and is characterized by the transient expansion of resident fibroblasts and the formation of a provisional matrix, as well as proliferation of airway progenitor cells and type II alveolar epithelial cells (AECII), with differentiation into type I alveolar epithelial cells (AECI).<sup>28</sup> During the fibrotic phase of ARDS (Panel B), which is strongly associated with the need for mechanical ventilation, extensive basement membrane damage and inadequate or delayed reepithelialization lead to the development of interstitial and intraalveolar fibrosis. AQP5 denotes aquaporin 5, CFTR cystic fibrosis transmembrane conductance regulator, GM-CSF granulocyte–macrophage colony-stimulating factor, HGF hepatocyte growth factor, IGF-I insulin-like growth factor I, IRF4 interferon regulatory factor 4, KGF keratinocyte growth factor, MR mannose receptor, PDGF platelet-derived growth factor, and TGF- $\beta$  transforming growth factor  $\beta$ .

if plateau airway pressures exceed 30 cm of water. In the landmark ARDS Network trial,<sup>33</sup> this approach, as compared with an approach involv-

ing a higher tidal volume, which had been used for decades, resulted in an absolute reduction of 9 percentage points in mortality. The respiratory

rate set on the ventilator may be increased to maintain acceptable minute ventilation (the volume of gas exhaled per minute) and carbon dioxide removal. However, recent preclinical and observational studies of mechanical power and energy transfer to the lung (proportional to lung elastance, tidal volume, pulmonary resistance, and respiratory rate) support prospective examination of a strategy using a lower respiratory rate.<sup>34</sup>

A PEEP of at least 5 cm of water is recommended, and a patient-level meta-analysis of three randomized trials suggests that mortality is increased when the PEEP is kept relatively low, as compared with a strategy involving a higher PEEP (a mean initial PEEP of approximately 16 cm of water), in patients with moderate-to-severe ARDS.<sup>32</sup> The optimal method for PEEP adjustment is unclear.<sup>35</sup> End-expiratory pleural pressure is often positive during ARDS (especially in patients with high abdominal pressures or obesity) and may be higher than traditionally applied levels of PEEP. This results in negative transpulmonary pressures at end-expiration, leading to atelectrauma. Measuring esophageal pressure with a manometer to estimate pleural pressure allows for adjustment of PEEP to achieve a positive end-expiratory transpulmonary pressure gradient, an approach that is increasingly used in clinical care. A small proof-of-concept study hinted at a reduction in mortality with the use of this strategy,<sup>36</sup> and clinical trials are under way (ClinicalTrials.gov numbers, NCT01681225 and NCT02416037). Adjusting PEEP or tidal volume to minimize driving pressure (the difference between plateau airway pressure and PEEP) is also rational, since with this approach, the tidal volume is adjusted in proportion to the patient's respiratory system compliance (to avoid overdistention). Adjusting the PEEP to minimize driving pressure may align the PEEP with the best respiratory system compliance, thus balancing the opening of the lung (and preventing atelectrauma) against overdistention (limiting volutrauma).<sup>37</sup>

In cases of moderate-to-severe ARDS (ratio of the partial pressure of arterial oxygen [ $P_{aO_2}$ ] to the fraction of inspired oxygen [ $F_{iO_2}$ ],  $<120$  mm Hg), ventilation while the patient is in the prone position is associated with reduced mortality and is currently recommended.<sup>30,32,38</sup> A benefit is likely to accrue from reducing the risk of ventilator-associated lung injury through the combined effects of more uniform distribution of ventila-

tion and less compression of the left lower lobe (by the heart).<sup>39</sup> As compared with deep sedation alone, neuromuscular blockade has been shown to improve outcomes in patients with moderate-to-severe ARDS ( $P_{aO_2}:F_{iO_2}$ ,  $<150$  mm Hg), possibly because neuromuscular blockage ensures patient-ventilator synchrony, which in turn reduces the risk of ventilator-associated lung injury.<sup>40</sup> However, deep sedation may be associated with deleterious effects on its own. Consequently, another large, randomized clinical trial involving patients with moderate-to-severe ARDS is comparing neuromuscular blockade and deep sedation with no routine neuromuscular blockade and less sedation (NCT02509078).<sup>41</sup>

High-frequency oscillation offers no advantage over conventional ventilation strategies and may be harmful, though a patient-level meta-analysis has suggested a benefit when the  $P_{aO_2}:F_{iO_2}$  ratio is less than 60 mm Hg.<sup>42-44</sup> Airway pressure release ventilation (i.e., applied continuous positive airway pressure that at a set interval releases the applied pressure) may improve oxygenation and tolerance of mechanical ventilation but has not been proved to reduce mortality. Both these ventilation strategies may improve oxygenation by increasing the mean airway pressure, which may adversely affect hemodynamics. Extracorporeal membrane oxygenation (ECMO) is reserved for patients with very severe ARDS ( $P_{aO_2}:F_{iO_2}$ ,  $<60$  mm Hg) after adequate lung-protective practices and correction of volume overload have failed to improve oxygenation. One randomized trial suggested a benefit with referral to an ECMO center, although it is unclear whether the benefit was simply from better specialized care, since not all referred patients were treated with ECMO.<sup>45</sup> A multicenter, randomized trial to further test the benefits of ECMO is ongoing (NCT01470703).

Noninvasive ventilation may increase the risk of death when attempted in patients with severe hypoxemia, perhaps by facilitating high tidal volumes from the combined effects of the high respiratory drive and respiratory support (resulting in ventilation-induced lung injury).<sup>46,47</sup> Oxygen administration through high-flow nasal cannulae and noninvasive ventilation provided with a helmet may be effective alternatives to intubation and mechanical ventilation in patients with less severe ARDS. Both approaches have the potential to reduce respiratory drive and the risk of ventilation-induced lung injury.<sup>47</sup>



A conservative fluid-management strategy shortened the duration of assisted ventilation in a large randomized trial,<sup>48</sup> and the benefit appears to occur largely from avoidance of fluid administration after reversal of shock.<sup>49</sup> Small randomized trials of diuretics and albumin after shock reversal showed improved oxygenation and a trend toward a shorter duration of mechanical ventilation,<sup>50,51</sup> but a larger trial did not suggest a reduction in mortality with the use of albumin in a general ICU population.<sup>52</sup> Albumin may be harmful in patients with traumatic brain injury.<sup>53</sup> For nutritional support, trophic and early full-calorie enteral nutrition are equivalent with regard to mortality,<sup>54</sup> and aggressive early caloric supplementation with parenteral nutrition may be harmful.<sup>55</sup>

#### PHARMACOLOGIC THERAPY

Unfortunately, no pharmacologic therapy for ARDS has been shown to reduce either short-term or long-term mortality. Inhaled nitric oxide transiently improves oxygenation and may improve long-term lung function among patients who survive, but it does not reduce mortality and is associated with acute kidney injury.<sup>56</sup> Glucocorticoids may improve oxygenation and airway pressures and, in patients with pneumonia, may hasten radiographic improvement, but these agents are not associated with a consistent survival benefit and are harmful if started 14 days or more after ARDS has been diagnosed.<sup>57</sup> Surfactant replacement, neutrophil elastase inhibition, and anticoagulation have failed in clinical trials, as have nonsteroidal antiinflammatory agents (ketoconazole and lysofylline), statins, albuterol, and antioxidants (procysteine [L-2-oxothiazolidine-4-carboxylic acid]), though many of these trials had relatively small samples, and in some cases, the doses tested did not modulate the intended biologic targets.<sup>58</sup> A trial of nebulized heparin is under way after promising early-phase testing (Australian New Zealand Clinical Trials Registry number, ACTRN12612000418875). A novel therapeutic approach in early clinical development involves intravenous delivery of mesenchymal stem cells, which interact with injured tissue through the release of multiple soluble bioactive factors.<sup>59</sup>

#### PREVENTION

With regard to prevention, observational studies indicate that a bundle of good ICU practices,

such as lower tidal volumes for all mechanically ventilated patients, early volume resuscitation and antibiotics for sepsis, male-donor plasma and restrictive use of blood products (to reduce the risk of transfusion-associated lung injury and volume overload), and intensivist involvement prevent the development of nosocomial ARDS.<sup>17</sup> Patients at risk for ARDS can be identified prospectively, allowing for trials of prevention and early treatment. The National Heart, Lung, and Blood Institute has funded a clinical trials network for this purpose. Thus far, glucocorticoids, aspirin, and beta-agonists have failed in prevention trials, although inhaled beta-agonists prevent high-altitude pulmonary edema and, in one small pilot trial, the combination of beta-agonists and glucocorticoids prevented the development of ARDS (but did not reduce mortality).<sup>17</sup>

#### THE SEARCH FOR TREATMENT-RESPONSIVE SUBTYPES

Patients at the more severe end of the ARDS spectrum have, on average, greater lung weights and higher rates of diffuse alveolar damage and pneumonia on biopsy or postmortem examination, and such patients are more likely to die from refractory hypoxemia than from multiorgan failure, which is the most common cause of death in all patients with ARDS.<sup>60</sup> As noted above, the subset of patients with severe ARDS appears to derive a survival benefit from treatments aimed at preventing ventilator-associated lung injury, including ventilation in the prone position and higher PEEP, two interventions that failed in unselected ARDS populations. However, lung histologic features are quite variable in all the Berlin definition subgroups,<sup>10</sup> including severe ARDS, and it is likely that distinct molecular mechanisms are involved.<sup>61,62</sup> A more precise medical approach will probably be necessary to identify pharmacotherapies for ARDS. For example, by merging clinical and biologic variables with genetic features, subphenotypes within asthma have been consistently identified, linked to specific molecular pathways, and shown to be responsive to different treatments.<sup>63</sup> This transformative approach to asthma is now unfolding for ARDS.

The most promising of these approaches involves latent class analysis of baseline clinical, laboratory, and protein biomarker levels, which has consistently identified a subpopulation of

patients with ARDS enrolled in clinical trials.<sup>61,62</sup> These patients, representing approximately one third of all patients with ARDS, have a “hyper-inflammatory” subphenotype, with elevated plasma levels of interleukin-6, interleukin-8, and tumor necrosis factor  $\alpha$  and reduced levels of bicarbonate and protein C. Sepsis and vasopressor use have been more common in this subpopulation than in other patients with ARDS, and mortality has been nearly twice as high, with randomized assignment to higher PEEP or conservative fluid management associated with a reduction in mortality.<sup>61,62</sup> Classification of the severity of ARDS and use of traditional clinical variables or severity-of-illness scores, such as the Acute Physiology, Age, and Chronic Health Evaluation (APACHE) III score, could not identify patients with this treatment-responsive subtype, but a relatively simple assessment of three to five biomarkers could. The biologic processes driv-

ing class assignment are unclear, and these promising findings require prospective validation. Additional approaches to the identification of treatment targets and responsive subtypes are described in the Supplementary Appendix.

## CONCLUSIONS

We now recognize that ARDS, like asthma, is a syndrome characterized by substantial heterogeneity.<sup>63</sup> A much better understanding of the biologic and genetic underpinnings of the subphenotypes of ARDS should lead the way to more targeted therapies. Until then, ICU practices that prevent ARDS, early and effective treatment of the insults leading to ARDS, and lung-protective ventilation and sensible fluid management remain the essential elements for good outcomes.

Disclosure forms provided by the authors are available with the full text of this article at NEJM.org.

## REFERENCES

- Ashbaugh DG, Bigelow DB, Petty TL, Levine BE. Acute respiratory distress in adults. *Lancet* 1967;2:319-23.
- Ware LB, Matthay MA. The acute respiratory distress syndrome. *N Engl J Med* 2000;342:1334-49.
- Bellani G, Laffey JG, Pham T, et al. Epidemiology, patterns of care, and mortality for patients with acute respiratory distress syndrome in intensive care units in 50 countries. *JAMA* 2016;315:788-800.
- Herridge MS, Moss M, Hough CL, et al. Recovery and outcomes after the acute respiratory distress syndrome (ARDS) in patients and their family caregivers. *Intensive Care Med* 2016;42:725-38.
- Herridge MS, Tansey CM, Matté A, et al. Functional disability 5 years after acute respiratory distress syndrome. *N Engl J Med* 2011;364:1293-304.
- Ferguson ND, Fan E, Camporota L, et al. The Berlin definition of ARDS: an expanded rationale, justification, and supplementary material. *Intensive Care Med* 2012;38:1573-82.
- Bellani G, Messa C, Guerra L, et al. Lungs of patients with acute respiratory distress syndrome show diffuse inflammation in normally aerated regions: a [18F]-fluoro-2-deoxy-D-glucose PET/CT study. *Crit Care Med* 2009;37:2216-22.
- Aublanc M, Perinel S, Guérin C. Acute respiratory distress syndrome mimics: the role of lung biopsy. *Curr Opin Crit Care* 2017;23:24-9.
- Katzenstein AL, Bloor CM, Leibow AA. Diffuse alveolar damage — the role of oxygen, shock, and related factors: a review. *Am J Pathol* 1976;85:209-28.
- Bernard GR, Artigas A, Brigham KL, et al. The American-European Consensus Conference on ARDS: definitions, mechanisms, relevant outcomes, and clinical trial coordination. *Am J Respir Crit Care Med* 1994;149:818-24.
- Esteban A, Fernández-Segoviano P, Frutos-Vivar F, et al. Comparison of clinical criteria for the acute respiratory distress syndrome with autopsy findings. *Ann Intern Med* 2004;141:440-5.
- Thille AW, Esteban A, Fernández-Segoviano P, et al. Comparison of the Berlin definition for acute respiratory distress syndrome with autopsy. *Am J Respir Crit Care Med* 2013;187:761-7.
- Villar J, Blanco J, Kacmarek RM. Current incidence and outcome of the acute respiratory distress syndrome. *Curr Opin Crit Care* 2016;22:1-6.
- Riviello ED, Kiviri W, Twagirumugabe T, et al. Hospital incidence and outcomes of the acute respiratory distress syndrome using the Kigali modification of the Berlin definition. *Am J Respir Crit Care Med* 2016;193:52-9.
- Eisner MD, Thompson T, Hudson LD, et al. Efficacy of low tidal volume ventilation in patients with different clinical risk factors for acute lung injury and the acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2001;164:231-6.
- Herasevich V, Yilmaz M, Khan H, Hubmayr RD, Gajic O. Validation of an electronic surveillance system for acute lung injury. *Intensive Care Med* 2009;35:1018-23.
- Yadav H, Thompson BT, Gajic O. Fifty years of research in ARDS: is acute respiratory distress syndrome a preventable disease? *Am J Respir Crit Care Med* 2017;195:725-36.
- Meyer NJ, Christie JD. Genetic heterogeneity and risk of acute respiratory distress syndrome. *Semin Respir Crit Care Med* 2013;34:459-74.
- Christie JD, Wurfel MM, Feng R, et al. Genome wide association identifies PPF1A1 as a candidate gene for acute lung injury risk following major trauma. *PLoS One* 2012;7(1):e28268.
- Marshall RP, Webb S, Bellingan GJ, et al. Angiotensin converting enzyme insertion/deletion polymorphism is associated with susceptibility and outcome in acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2002;166:646-50.
- Li W, Moore MJ, Vasileva N, et al. Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. *Nature* 2003;426:450-4.
- Kuba K, Imai Y, Rao S, et al. A crucial role of angiotensin converting enzyme 2 (ACE2) in SARS coronavirus-induced lung injury. *Nat Med* 2005;11:875-9.
- Imai Y, Kuba K, Rao S, et al. Angiotensin-converting enzyme 2 protects from severe acute lung failure. *Nature* 2005;436:112-6.
- Shortt K, Chaudhary S, Grigoryev D, et al. Identification of novel single nucleotide polymorphisms associated with acute respiratory distress syndrome by exome-seq. *PLoS One* 2014;9(11):e111953.
- Meyer NJ, Calfee CS. Novel translational approaches to the search for precision therapies for acute respiratory distress syndrome. *Lancet Respir Med* 2017;5:512-23.
- Huh D, Leslie DC, Matthews BD, et al. A human disease model of drug toxicity-induced pulmonary edema in a lung-on-a-chip microdevice. *Sci Transl Med* 2012;4:159ra147.

27. Lee JW, Krasnodemskaia A, McKenna DH, Song Y, Abbott J, Matthay MA. Therapeutic effects of human mesenchymal stem cells in ex vivo human lungs injured with live bacteria. *Am J Respir Crit Care Med* 2013;187:751-60.
28. Vaughan AE, Brumwell AN, Xi Y, et al. Lineage-negative progenitors mobilize to regenerate lung epithelium after major injury. *Nature* 2015;517:621-5.
29. Aggarwal NR, King LS, D'Alessio FR. Diverse macrophage populations mediate acute lung inflammation and resolution. *Am J Physiol Lung Cell Mol Physiol* 2014;306:L709-L725.
30. Rhodes A, Evans LE, Alhazzani W, et al. Surviving Sepsis Campaign: international guidelines for management of sepsis and septic shock: 2016. *Crit Care Med* 2017;45:486-552.
31. Slutsky AS, Ranieri VM. Ventilator-induced lung injury. *N Engl J Med* 2013;369:2126-36.
32. Fan E, Del Sorbo L, Goligher EC, et al. An official American Thoracic Society/European Society of Intensive Care Medicine/Society of Critical Care Medicine clinical practice guideline: mechanical ventilation in adult patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2017;195:1253-63.
33. The Acute Respiratory Distress Syndrome Network. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med* 2000;342:1301-8.
34. Gattinoni L, Tonetti T, Cressoni M, et al. Ventilator-related causes of lung injury: the mechanical power. *Intensive Care Med* 2016;42:1567-75.
35. Briel M, Meade M, Mercat A, et al. Higher vs lower positive end-expiratory pressure in patients with acute lung injury and acute respiratory distress syndrome: systematic review and meta-analysis. *JAMA* 2010;303:865-73.
36. Talmor D, Sarge T, Malhotra A, et al. Mechanical ventilation guided by esophageal pressure in acute lung injury. *N Engl J Med* 2008;359:2095-104.
37. Amato MBP, Meade MO, Slutsky AS, et al. Driving pressure and survival in the acute respiratory distress syndrome. *N Engl J Med* 2015;372:747-55.
38. Guérin C, Reignier J, Richard J-C, et al. Prone positioning in severe acute respiratory distress syndrome. *N Engl J Med* 2013;368:2159-68.
39. Gattinoni L, Tognoni G, Pesenti A, et al. Effect of prone positioning on the survival of patients with acute respiratory failure. *N Engl J Med* 2001;345:568-73.
40. Papazian L, Forel J-M, Gacouin A, et al. Neuromuscular blockers in early acute respiratory distress syndrome. *N Engl J Med* 2010;363:1107-16.
41. Huang DT, Angus DC, Moss M, et al. Design and rationale of the Reevaluation of Systemic Early Neuromuscular Blockade Trial for acute respiratory distress syndrome. *Ann Am Thorac Soc* 2017;14:124-33.
42. Young D, Lamb SE, Shah S, et al. High-frequency oscillation for acute respiratory distress syndrome. *N Engl J Med* 2013;368:806-13.
43. Ferguson ND, Cook DJ, Guyatt GH, et al. High-frequency oscillation in early acute respiratory distress syndrome. *N Engl J Med* 2013;368:795-805.
44. Meade MO, Young D, Hanna S, et al. Severity of hypoxemia and effect of high frequency oscillatory ventilation in ARDS. *Am J Respir Crit Care Med* 2017 February 28 (Epub ahead of print).
45. Noah MA, Peek GJ, Finney SJ, et al. Referral to an extracorporeal membrane oxygenation center and mortality among patients with severe 2009 influenza A(H1N1). *JAMA* 2011;306:1659-68.
46. Bellani G, Laffey JG, Pham T, et al. Noninvasive ventilation of patients with acute respiratory distress syndrome: insights from the LUNG SAFE study. *Am J Respir Crit Care Med* 2017;195:67-77.
47. Brochard L, Slutsky A, Pesenti A. Mechanical ventilation to minimize progression of lung injury in acute respiratory failure. *Am J Respir Crit Care Med* 2017;195:438-42.
48. The National Heart, Lung, and Blood Institute Acute Respiratory Distress Syndrome (ARDS) Clinical Trials Network. Comparison of two fluid-management strategies in acute lung injury. *N Engl J Med* 2006;354:2564-75.
49. Semler MW, Wheeler AP, Thompson BT, Bernard GR, Wiedemann HP, Rice TW. Impact of initial central venous pressure on outcomes of conservative versus liberal fluid management in acute respiratory distress syndrome. *Crit Care Med* 2016;44:782-9.
50. Martin GS, Moss M, Wheeler AP, Mealer M, Morris JA, Bernard GR. A randomized, controlled trial of furosemide with or without albumin in hypoproteinemic patients with acute lung injury. *Crit Care Med* 2005;33:1681-7.
51. Martin GS, Mangialardi RJ, Wheeler AP, Dupont WD, Morris JA, Bernard GR. Albumin and furosemide therapy in hypoproteinemic patients with acute lung injury. *Crit Care Med* 2002;30:2175-82.
52. The SAFE Study Investigators. A comparison of albumin and saline for fluid resuscitation in the intensive care unit. *N Engl J Med* 2004;350:2247-56.
53. The SAFE Study Investigators. Saline or albumin for fluid resuscitation in patients with traumatic brain injury. *N Engl J Med* 2007;357:874-84.
54. The National Heart, Lung, and Blood Institute Acute Respiratory Distress Syndrome (ARDS) Clinical Trials Network. Initial trophic vs full enteral feeding in patients with acute lung injury: the EDEN randomized trial. *JAMA* 2012;307:795-803.
55. Casaer MP, Mesotten D, Hermans G, et al. Early versus late parenteral nutrition in critically ill adults. *N Engl J Med* 2011;365:506-17.
56. Griffiths MJ, Evans TW. Inhaled nitric oxide therapy in adults. *N Engl J Med* 2005;353:2683-95.
57. The National Heart, Lung, and Blood Institute Acute Respiratory Distress Syndrome (ARDS) Clinical Trials Network. Efficacy and safety of corticosteroids for persistent acute respiratory distress syndrome. *N Engl J Med* 2006;354:1671-84.
58. Boyle AJ, Mac Sweeney R, McAuley DF. Pharmacological treatments in ARDS; a state-of-the-art update. *BMC Med* 2013;11:166.
59. Wilson JG, Liu KD, Zhuo H, et al. Mesenchymal stem (stromal) cells for treatment of ARDS: a phase 1 clinical trial. *Lancet Respir Med* 2015;3:24-32.
60. Thompson BT, Guérin C, Esteban A. Should ARDS be renamed diffuse alveolar damage? *Intensive Care Med* 2016;42:653-5.
61. Calfee CS, Delucchi K, Parsons PE, Thompson BT, Ware LB, Matthay MA. Subphenotypes in acute respiratory distress syndrome: latent class analysis of data from two randomised controlled trials. *Lancet Respir Med* 2014;2:611-20.
62. Famous KR, Delucchi K, Ware LB, et al. Acute respiratory distress syndrome subphenotypes respond differently to randomized fluid management strategy. *Am J Respir Crit Care Med* 2017;195:331-8.
63. Wenzel SE. Asthma phenotypes: the evolution from clinical to molecular approaches. *Nat Med* 2012;18:716-25.

Copyright © 2017 Massachusetts Medical Society.